

Water Indicators

Indicator	Value	Description	Source
Overall Basin Risk (score)	3.02	Overall Basin Risk (score)	
Overall Basin Risk (rank)	27	Overall Basin Risk (rank)	
Physical risk (score)	3.01	Physical risk (score)	
Physical risk (rank)	49	Physical risk (rank)	
Regulatory risk (score)	2.91	Regulatory risk (score)	
Regulatory risk (rank)	81	Regulatory risk (rank)	
Reputation risk (score)	3.14	Reputation risk (score)	
Reputation risk (rank)	41	Reputation risk (rank)	
1. Quantity - Scarcity (score)	4.23	1. Quantity - Scarcity (score)	
1. Quantity - Scarcity (rank)	7	1. Quantity - Scarcity (rank)	
2. Quantity - Flooding (score)	1.82	2. Quantity - Flooding (score)	
2. Quantity - Flooding (rank)	171	2. Quantity - Flooding (rank)	
3. Quality (score)	2.00	3. Quality (score)	
3. Quality (rank)	149	3. Quality (rank)	
4. Ecosystem Service Status (score)	1.65	4. Ecosystem Service Status (score)	
4. Ecosystem Service Status (rank)	163	4. Ecosystem Service Status (rank)	
5. Enabling Environment (Policy & Laws) (score)	2.55	5. Enabling Environment (Policy & Laws) (score)	
5. Enabling Environment (Policy & Laws) (rank)	110	5. Enabling Environment (Policy & Laws) (rank)	
6. Institutions and Governance (score)	4.25	6. Institutions and Governance (score)	
6. Institutions and Governance (rank)	18	6. Institutions and Governance (rank)	
7. Management Instruments (score)	2.44	7. Management Instruments (score)	
7. Management Instruments (rank)	125	7. Management Instruments (rank)	
8 - Infrastructure & Finance (score)	1.75	8 - Infrastructure & Finance (score)	
8 - Infrastructure & Finance (rank)	109	8 - Infrastructure & Finance (rank)	
9. Cultural Diversity (score)	1.00	9. Cultural importance (score)	
9. Cultural Diversity (rank)	129	9. Cultural importance (rank)	
10. Biodiversity Importance (score)	1.86	10. Biodiversity importance (score)	

Country Overview - Egypt

Indicator	Value	Description	Source
10. Biodiversity Importance (rank)	186	10. Biodiversity importance (rank)	
11. Media Scrutiny (score)	4.00	11. Media Scrutiny (score)	
11. Media Scrutiny (rank)	11	11. Media Scrutiny (rank)	
12. Conflict (score)	3.53	12. Conflict (score)	
12. Conflict (rank)	13	12. Conflict (rank)	
1.0 - Aridity (score)	4.86	The aridity risk indicator is based on the Global Aridity Index (Global-Aridity) and Global Potential Evapo-Transpiration (Global-PET) Geospatial data sets by Trabucco and Zomer (2009). These data sets provide information about the potential availability of water in regions with low water demand, thus they are used in the Water Risk Filter 5.0 to better account for deserts and other arid areas in the risk assessment.	Trabucco, A., & Zomer, R. J. (2009). Global potential evapo-transpiration (Global-PET) and global aridity index (Global-Aridity) geodatabase. CGIAR consortium for spatial information.
1.0 - Aridity (rank)	1	The aridity risk indicator is based on the Global Aridity Index (Global-Aridity) and Global Potential Evapo-Transpiration (Global-PET) Geospatial data sets by Trabucco and Zomer (2009). These data sets provide information about the potential availability of water in regions with low water demand, thus they are used in the Water Risk Filter 5.0 to better account for deserts and other arid areas in the risk assessment.	Trabucco, A., & Zomer, R. J. (2009). Global potential evapo-transpiration (Global-PET) and global aridity index (Global-Aridity) geodatabase. CGIAR consortium for spatial information.
1.1 - Water Depletion (score)	3.84	The water depletion risk indicator is based on annual average monthly net water depletion from Brauman et al. (2016). Their analysis is based on model outputs from the newest version of the integrated water resources model WaterGAP3 which measures water depletion as the ratio of water consumption-to-availability.	Brauman, K. A., Richter, B. D., Postel, S., Malsy, M., & Flörke, M. (2016). Water depletion: An improved metric for incorporating seasonal and dry-year water scarcity into water risk assessments. <i>Elem Sci Anth</i> , 4.
1.1 - Water Depletion (rank)	9	The water depletion risk indicator is based on annual average monthly net water depletion from Brauman et al. (2016). Their analysis is based on model outputs from the newest version of the integrated water resources model WaterGAP3 which measures water depletion as the ratio of water consumption-to-availability.	Brauman, K. A., Richter, B. D., Postel, S., Malsy, M., & Flörke, M. (2016). Water depletion: An improved metric for incorporating seasonal and dry-year water scarcity into water risk assessments. <i>Elem Sci Anth</i> , 4.
1.2 - Baseline Water Stress (score)	3.63	World Resources Institute's Baseline Water Stress measures the ratio of total annual water withdrawals to total available annual renewable supply, accounting for upstream consumptive use. A higher percentage indicates more competition among users.	Hofste, R., Kuzma, S., Walker, S., ... & Sutanudjaja, E.H. (2019). <i>Aqueduct 3.0: Updated decision relevant global water risk indicators</i> . Technical note. Washington, DC: World Resources Institute.

Country Overview - Egypt

Indicator	Value	Description	Source
1.2 - Baseline Water Stress (rank)	32	World Resources Institute's Baseline Water Stress measures the ratio of total annual water withdrawals to total available annual renewable supply, accounting for upstream consumptive use. A higher percentage indicates more competition among users.	Hofste, R., Kuzma, S., Walker, S., ... & Sutanudjaja, E.H. (2019). Aqueduct 3.0: Updated decision relevant global water risk indicators. Technical note. Washington, DC: World Resources Institute.
1.3 - Blue Water Scarcity (score)	4.92	The blue water scarcity risk indicator is based on Mekonnen and Hoekstra (2016) global assessment of blue water scarcity on a monthly basis and at high spatial resolution (grid cells of 30 × 30 arc min resolution). Blue water scarcity is calculated as the ratio of the blue water footprint in a grid cell to the total blue water availability in the cell. The time period analyzed in this study ranges from 1996 to 2005.	Mekonnen, M. M., & Hoekstra, A. Y. (2016). Four billion people facing severe water scarcity. <i>Science advances</i> , 2(2), e1500323.
1.3 - Blue Water Scarcity (rank)	12	The blue water scarcity risk indicator is based on Mekonnen and Hoekstra (2016) global assessment of blue water scarcity on a monthly basis and at high spatial resolution (grid cells of 30 × 30 arc min resolution). Blue water scarcity is calculated as the ratio of the blue water footprint in a grid cell to the total blue water availability in the cell. The time period analyzed in this study ranges from 1996 to 2005.	Mekonnen, M. M., & Hoekstra, A. Y. (2016). Four billion people facing severe water scarcity. <i>Science advances</i> , 2(2), e1500323.
1.4 - Projected Change in Water Discharge (by ~2050) (score)	2.11	This risk indicator is based on multi-model simulation that applies both global climate and hydrological models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP). To estimate the change at 2°C of global warming above 1980-2010 levels, simulated annual water discharge was averaged over a 31-year period with 2°C mean warming. Results are expressed in terms of relative change (%) in probability between present day (1980-2010) conditions and 2°C scenarios by 2050.	Schewe, J., Heinke, J., Gerten, D., Haddeland, I., Arnell, N. W., Clark, D. B., ... & Gosling, S. N. (2014). Multimodel assessment of water scarcity under climate change. <i>Proceedings of the National Academy of Sciences</i> , 111(9), 3245-3250.
1.4 - Projected Change in Water Discharge (by ~2050) (rank)	72	This risk indicator is based on multi-model simulation that applies both global climate and hydrological models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP). To estimate the change at 2°C of global warming above 1980-2010 levels, simulated annual water discharge was averaged over a 31-year period with 2°C mean warming. Results are expressed in terms of relative change (%) in probability between present day (1980-2010) conditions and 2°C scenarios by 2050.	Schewe, J., Heinke, J., Gerten, D., Haddeland, I., Arnell, N. W., Clark, D. B., ... & Gosling, S. N. (2014). Multimodel assessment of water scarcity under climate change. <i>Proceedings of the National Academy of Sciences</i> , 111(9), 3245-3250.

Country Overview - Egypt

Indicator	Value	Description	Source
1.5 - Drought Frequency Probability (score)	5.00	This risk indicator is based on the Standardized Precipitation and Evaporation Index (SPEI). Vicente-Serrano et al. (2010) developed this multi-scalar drought index applying both precipitation and temperature data to detect, monitor and analyze different drought types and impacts in the context of global warming. The mathematical calculations used for SPEI are similar to the Standard Precipitation Index (SPI), but it has the advantage to include the role of evapotranspiration.	Vicente-Serrano, S. M., Beguería, S., & López-Moreno, J. I. (2010). A multiscale drought index sensitive to global warming: the standardized precipitation evapotranspiration index. <i>Journal of climate</i> , 23(7), 1696-1718.
1.5 - Drought Frequency Probability (rank)	4	This risk indicator is based on the Standardized Precipitation and Evaporation Index (SPEI). Vicente-Serrano et al. (2010) developed this multi-scalar drought index applying both precipitation and temperature data to detect, monitor and analyze different drought types and impacts in the context of global warming. The mathematical calculations used for SPEI are similar to the Standard Precipitation Index (SPI), but it has the advantage to include the role of evapotranspiration.	Vicente-Serrano, S. M., Beguería, S., & López-Moreno, J. I. (2010). A multiscale drought index sensitive to global warming: the standardized precipitation evapotranspiration index. <i>Journal of climate</i> , 23(7), 1696-1718.
1.6 - Projected Change in Drought Occurrence (by ~2050) (score)	3.12	This risk indicator is based on multi-model simulation that applies both global climate and drought models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP) . A drought threshold for pre-industrial conditions was calculated based on time-series averages. Results are expressed in terms of relative change (%) in probability between pre-industrial and 2°C scenarios.	Frieler, K., Lange, S., Piontek, F., Reyer, C. P., Schewe, J., Warszawski, L., ... & Geiger, T. (2017). Assessing the impacts of 1.5 C global warming-simulation protocol of the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP2b). Geoscientific Model Development.
1.6 - Projected Change in Drought Occurrence (by ~2050) (rank)	50	This risk indicator is based on multi-model simulation that applies both global climate and drought models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP) . A drought threshold for pre-industrial conditions was calculated based on time-series averages. Results are expressed in terms of relative change (%) in probability between pre-industrial and 2°C scenarios.	Frieler, K., Lange, S., Piontek, F., Reyer, C. P., Schewe, J., Warszawski, L., ... & Geiger, T. (2017). Assessing the impacts of 1.5 C global warming-simulation protocol of the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP2b). Geoscientific Model Development.
2.1 - Estimated Flood Occurrence (score)	1.74	This risk indicator is based on the recurrence of floods within the 34-year time frame period of 1985 to 2019. The occurrence of floods within a given location was estimated using data from Flood Observatory, University of Colorado. The Flood Observatory use data derived from a wide variety of news, governmental, instrumental, and remote sensing source.	Brakenridge, G. R. (2019). Global active archive of large flood events. Dartmouth Flood Observatory, University of Colorado.
2.1 - Estimated Flood Occurrence (rank)	174	This risk indicator is based on the recurrence of floods within the 34-year time frame period of 1985 to 2019. The occurrence of floods within a given location was estimated using data from Flood Observatory, University of Colorado. The Flood Observatory use data derived from a wide variety of news, governmental, instrumental, and remote sensing source.	Brakenridge, G. R. (2019). Global active archive of large flood events. Dartmouth Flood Observatory, University of Colorado.

Indicator	Value	Description	Source
2.2 - Projected Change in Flood Occurrence (by ~2050) (score)	3.28	This risk indicator is based on multi-model simulation that applies both global climate and drought models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP). The magnitude of the flood event was defined based on 100-year return period for pre-industrial conditions. Results are expressed in terms of change (%) in probability between pre-industrial and 2°C scenarios.	Frieler, K., Lange, S., Piontek, F., Reyer, C. P., Schewe, J., Warszawski, L., ... & Geiger, T. (2017). Assessing the impacts of 1.5 C global warming-simulation protocol of the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP2b). Geoscientific Model Development.
2.2 - Projected Change in Flood Occurrence (by ~2050) (rank)	15	This risk indicator is based on multi-model simulation that applies both global climate and drought models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP). The magnitude of the flood event was defined based on 100-year return period for pre-industrial conditions. Results are expressed in terms of change (%) in probability between pre-industrial and 2°C scenarios.	Frieler, K., Lange, S., Piontek, F., Reyer, C. P., Schewe, J., Warszawski, L., ... & Geiger, T. (2017). Assessing the impacts of 1.5 C global warming-simulation protocol of the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP2b). Geoscientific Model Development.
3.1 - Surface Water Contamination Index (score)	2.00	<p>The underlying data for this risk indicator is based on a broad suite of pollutants with well-documented direct or indirect negative effects on water security for both humans and freshwater biodiversity, compiled by Vörösmarty et al. (2010). The negative effects are specific to individual pollutants, ranging from impacts mediated by eutrophication such as algal blooms and oxygen depletion (e.g., caused by phosphorus and organic loading) to direct toxic effects (e.g., caused by pesticides, mercury).</p> <p>The overall Surface Water Contamination Index is calculated based on a range of key pollutants with different weightings according to the level of their negative effects on water security for both humans and freshwater biodiversity: soil salinization (8%), nitrogen (12%) and phosphorus (P, 13%) loading, mercury deposition (5%), pesticide loading (10%), sediment loading (17%), organic loading (as Biological Oxygen Demand, BOD; 15%), potential acidification (9%), and thermal alteration (11%).</p>	Vörösmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P., ... & Davies, P. M. (2010). Global threats to human water security and river biodiversity. <i>Nature</i> , 467(7315), 555.

Indicator	Value	Description	Source
3.1 - Surface Water Contamination Index (rank)	149	<p>The underlying data for this risk indicator is based on a broad suite of pollutants with well-documented direct or indirect negative effects on water security for both humans and freshwater biodiversity, compiled by Vörösmarty et al. (2010). The negative effects are specific to individual pollutants, ranging from impacts mediated by eutrophication such as algal blooms and oxygen depletion (e.g., caused by phosphorus and organic loading) to direct toxic effects (e.g., caused by pesticides, mercury).</p> <p>The overall Surface Water Contamination Index is calculated based on a range of key pollutants with different weightings according to the level of their negative effects on water security for both humans and freshwater biodiversity: soil salinization (8%), nitrogen (12%) and phosphorus (P, 13%) loading, mercury deposition (5%), pesticide loading (10%), sediment loading (17%), organic loading (as Biological Oxygen Demand, BOD; 15%), potential acidification (9%), and thermal alteration (11%).</p>	Vörösmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P., ... & Davies, P. M. (2010). Global threats to human water security and river biodiversity. <i>Nature</i> , 467(7315), 555.
4.1 - Fragmentation Status of Rivers (score)	1.89	<p>This risk indicator is based on the data set by Grill et al. (2019) mapping the world's free-flowing rivers. Grill et al. (2019) compiled a geometric network of the global river system and associated attributes, such as hydro-geometric properties, as well as pressure indicators to calculate an integrated connectivity status index (CSI). While only rivers with high levels of connectivity in their entire length are classified as free-flowing, rivers of CSI < 95% are considered as fragmented at a certain degree.</p>	Grill, G., Lehner, B., Thieme, M., Geenen, B., Tickner, D., Antonelli, F., ... & Macedo, H. E. (2019). Mapping the world's free-flowing rivers. <i>Nature</i> , 569(7755), 215.
4.1 - Fragmentation Status of Rivers (rank)	137	<p>This risk indicator is based on the data set by Grill et al. (2019) mapping the world's free-flowing rivers. Grill et al. (2019) compiled a geometric network of the global river system and associated attributes, such as hydro-geometric properties, as well as pressure indicators to calculate an integrated connectivity status index (CSI). While only rivers with high levels of connectivity in their entire length are classified as free-flowing, rivers of CSI < 95% are considered as fragmented at a certain degree.</p>	Grill, G., Lehner, B., Thieme, M., Geenen, B., Tickner, D., Antonelli, F., ... & Macedo, H. E. (2019). Mapping the world's free-flowing rivers. <i>Nature</i> , 569(7755), 215.
4.2 - Catchment Ecosystem Services Degradation Level (tree cover loss) (score)	1.00	<p>For this risk indicator, tree cover loss was applied as a proxy to represent catchment ecosystem services degradation since forests play an important role in terms of water regulation, supply and pollution control.</p> <p>The forest cover data is based on Hansen et al.'s global Landsat data at a 30-meter spatial resolution to characterize forest cover and change. The authors defined trees as vegetation taller than 5 meters in height, and forest cover loss as a stand-replacement disturbance, or a change from a forest to non-forest state, during the period 2000 – 2018.</p>	Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A. A., Tyukavina, A., ... & Kommareddy, A. (2013). High-resolution global maps of 21st-century forest cover change. <i>science</i> , 342(6160), 850-853.

Country Overview - Egypt

Indicator	Value	Description	Source
4.2 - Catchment Ecosystem Services Degradation Level (tree cover loss) (rank)	159	<p>For this risk indicator, tree cover loss was applied as a proxy to represent catchment ecosystem services degradation since forests play an important role in terms of water regulation, supply and pollution control.</p> <p>The forest cover data is based on Hansen et al.'s global Landsat data at a 30-meter spatial resolution to characterize forest cover and change. The authors defined trees as vegetation taller than 5 meters in height, and forest cover loss as a stand-replacement disturbance, or a change from a forest to non-forest state, during the period 2000 – 2018.</p>	Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A. A., Tyukavina, A., ... & Kommareddy, A. (2013). High-resolution global maps of 21st-century forest cover change. <i>science</i> , 342(6160), 850-853.
4.3 - Projected Impacts on Freshwater Biodiversity (score)	1.64	The study by Tedesco et al. (2013) to project changes [% increase or decrease] in extinction rate by ~2090 of freshwater fish due to water availability loss from climate change is used as a proxy to estimate the projected impacts on freshwater biodiversity.	Tedesco, P. A., Oberdorff, T., Cornu, J. F., Beauchard, O., Brosse, S., Dürr, H. H., ... & Hugueny, B. (2013). A scenario for impacts of water availability loss due to climate change on riverine fish extinction rates. <i>Journal of Applied Ecology</i> , 50(5), 1105-1115.
4.3 - Projected Impacts on Freshwater Biodiversity (rank)	165	The study by Tedesco et al. (2013) to project changes [% increase or decrease] in extinction rate by ~2090 of freshwater fish due to water availability loss from climate change is used as a proxy to estimate the projected impacts on freshwater biodiversity.	Tedesco, P. A., Oberdorff, T., Cornu, J. F., Beauchard, O., Brosse, S., Dürr, H. H., ... & Hugueny, B. (2013). A scenario for impacts of water availability loss due to climate change on riverine fish extinction rates. <i>Journal of Applied Ecology</i> , 50(5), 1105-1115.
5.1 - Freshwater Policy Status (SDG 6.5.1) (score)	2.00	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "National Water Resources Policy" indicator, which corresponds to one of the three national level indicators under the Enabling Environment category.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
5.1 - Freshwater Policy Status (SDG 6.5.1) (rank)	113	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "National Water Resources Policy" indicator, which corresponds to one of the three national level indicators under the Enabling Environment category.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
5.2 - Freshwater Law Status (SDG 6.5.1) (score)	3.00	<p>This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "National Water Resources Law(s)" indicator, which corresponds to one of the three national level indicators under the Enabling Environment category.</p> <p>For SDG 6.5.1, enabling environment depicts the conditions that help to support the implementation of IWRM, which includes legal and strategic planning tools for IWRM.</p>	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.

Country Overview - Egypt

Indicator	Value	Description	Source
5.2 - Freshwater Law Status (SDG 6.5.1) (rank)	54	<p>This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation “National Water Resources Law(s)” indicator, which corresponds to one of the three national level indicators under the Enabling Environment category.</p> <p>For SDG 6.5.1, enabling environment depicts the conditions that help to support the implementation of IWRM, which includes legal and strategic planning tools for IWRM.</p>	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
5.3 - Implementation Status of Water Management Plans (SDG 6.5.1) (score)	2.00	<p>This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation “National IWRM plans” indicator, which corresponds to one of the three national level indicators under the Enabling Environment category.</p> <p>For SDG 6.5.1, enabling environment depicts the conditions that help to support the implementation of IWRM, which includes legal and strategic planning tools for IWRM.</p>	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
5.3 - Implementation Status of Water Management Plans (SDG 6.5.1) (rank)	129	<p>This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation “National IWRM plans” indicator, which corresponds to one of the three national level indicators under the Enabling Environment category.</p> <p>For SDG 6.5.1, enabling environment depicts the conditions that help to support the implementation of IWRM, which includes legal and strategic planning tools for IWRM.</p>	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
6.1 - Corruption Perceptions Index (score)	4.00	<p>This risk Indicator is based on the latest Transparency International's data: the Corruption Perceptions Index 2018. This index aggregates data from a number of different sources that provide perceptions of business people and country experts on the level of corruption in the public sector.</p>	Transparency International (2019). Corruption Perceptions Index 2018. Berlin: Transparency International.
6.1 - Corruption Perceptions Index (rank)	31	<p>This risk Indicator is based on the latest Transparency International's data: the Corruption Perceptions Index 2018. This index aggregates data from a number of different sources that provide perceptions of business people and country experts on the level of corruption in the public sector.</p>	Transparency International (2019). Corruption Perceptions Index 2018. Berlin: Transparency International.
6.2 - Freedom in the World Index (score)	5.00	<p>This risk indicator is based on Freedom House (2019), an annual global report on political rights and civil liberties, composed of numerical ratings and descriptive texts for each country and a select group of territories. The 2019 edition involved more than 100 analysts and more than 30 advisers with global, regional, and issue-based expertise to covers developments in 195 countries and 14 territories from January 1, 2018, through December 31, 2018.</p>	Freedom House (2019). Freedom in the world 2019. Washington, DC: Freedom House.

Indicator	Value	Description	Source
6.2 - Freedom in the World Index (rank)	13	This risk indicator is based on Freedom House (2019), an annual global report on political rights and civil liberties, composed of numerical ratings and descriptive texts for each country and a select group of territories. The 2019 edition involved more than 100 analysts and more than 30 advisers with global, regional, and issue-based expertise to covers developments in 195 countries and 14 territories from January 1, 2018, through December 31, 2018.	Freedom House (2019). Freedom in the world 2019. Washington, DC: Freedom House.
6.3 - Business Participation in Water Management (SDG 6.5.1) (score)	4.00	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "Business Participation in Water Resources Development, Management and Use" indicator, which corresponds to one of the six national level indicators under the Institutions and Participation category.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
6.3 - Business Participation in Water Management (SDG 6.5.1) (rank)	11	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "Business Participation in Water Resources Development, Management and Use" indicator, which corresponds to one of the six national level indicators under the Institutions and Participation category.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
7.1 - Management Instruments for Water Management (SDG 6.5.1) (score)	2.00	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "Sustainable and efficient water use management" indicator, which corresponds to one of the five national level indicators under the Management Instruments category. For SDG 6.5.1, management instruments refer to the tools and activities that enable decision-makers and users to make rational and informed choices between alternative actions.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
7.1 - Management Instruments for Water Management (SDG 6.5.1) (rank)	119	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "Sustainable and efficient water use management" indicator, which corresponds to one of the five national level indicators under the Management Instruments category. For SDG 6.5.1, management instruments refer to the tools and activities that enable decision-makers and users to make rational and informed choices between alternative actions.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.

Country Overview - Egypt

Indicator	Value	Description	Source
7.2 - Groundwater Monitoring Data Availability and Management (score)	3.00	This risk indicator is based on the data set by UN IGRAC (2019) to determine the level of availability of groundwater monitoring data at country level as groundwater management decisions rely strongly on data availability. The level of groundwater monitoring data availability for groundwater management is determined according to a combination of three criteria developed by WWF and IGRAC: 1) Status of country groundwater monitoring programme, 2) groundwater data availability for NGOs and 3) Public access to processed groundwater monitoring data.	UN IGRAC (2019). Global Groundwater Monitoring Network GGMM Portal. UN International Groundwater Resources Assessment Centre (IGRAC).
7.2 - Groundwater Monitoring Data Availability and Management (rank)	42	This risk indicator is based on the data set by UN IGRAC (2019) to determine the level of availability of groundwater monitoring data at country level as groundwater management decisions rely strongly on data availability. The level of groundwater monitoring data availability for groundwater management is determined according to a combination of three criteria developed by WWF and IGRAC: 1) Status of country groundwater monitoring programme, 2) groundwater data availability for NGOs and 3) Public access to processed groundwater monitoring data.	UN IGRAC (2019). Global Groundwater Monitoring Network GGMM Portal. UN International Groundwater Resources Assessment Centre (IGRAC).
7.3 - Density of Runoff Monitoring Stations (score)	3.91	The density of monitoring stations for water quantity was applied as proxy to develop this risk indicator. The Global Runoff Data Base was used to estimate the number of monitoring stations per 1000km ² of the main river system (data base access date: May 2018).	BfG (2019). Global Runoff Data Base. German Federal Institute of Hydrology (BfG).
7.3 - Density of Runoff Monitoring Stations (rank)	59	The density of monitoring stations for water quantity was applied as proxy to develop this risk indicator. The Global Runoff Data Base was used to estimate the number of monitoring stations per 1000km ² of the main river system (data base access date: May 2018).	BfG (2019). Global Runoff Data Base. German Federal Institute of Hydrology (BfG).
8.1 - Access to Safe Drinking Water (score)	1.00	This risk indicator is based on the Joint Monitoring Programme for Water Supply, Sanitation and Hygiene (UNICEF/WHO) 2019 data. It provides estimates on the use of water, sanitation and hygiene by country for the period 2000-2017.	WHO & UNICEF (2019). Estimates on the use of water, sanitation and hygiene by country (2000-2017). Joint Monitoring Programme for Water Supply, Sanitation and Hygiene.
8.1 - Access to Safe Drinking Water (rank)	101	This risk indicator is based on the Joint Monitoring Programme for Water Supply, Sanitation and Hygiene (UNICEF/WHO) 2019 data. It provides estimates on the use of water, sanitation and hygiene by country for the period 2000-2017.	WHO & UNICEF (2019). Estimates on the use of water, sanitation and hygiene by country (2000-2017). Joint Monitoring Programme for Water Supply, Sanitation and Hygiene.
8.2 - Access to Sanitation (score)	2.00	This risk indicator is based on the Joint Monitoring Programme for Water Supply, Sanitation and Hygiene (UNICEF/WHO) 2019 data. It provides estimates on the use of water, sanitation and hygiene by country for the period 2000-2017.	WHO & UNICEF (2019). Estimates on the use of water, sanitation and hygiene by country (2000-2017). Joint Monitoring Programme for Water Supply, Sanitation and Hygiene.

Country Overview - Egypt

Indicator	Value	Description	Source
8.2 - Access to Sanitation (rank)	103	This risk indicator is based on the Joint Monitoring Programme for Water Supply, Sanitation and Hygiene (UNICEF/WHO) 2019 data. It provides estimates on the use of water, sanitation and hygiene by country for the period 2000-2017.	WHO & UNICEF (2019). Estimates on the use of water, sanitation and hygiene by country (2000-2017). Joint Monitoring Programme for Water Supply, Sanitation and Hygiene.
8.3 - Financing for Water Resource Development and Management (SDG 6.5.1) (score)	4.00	This risk indicator is based on the average 'Financing' score of UN SDG 6.5.1. Degree of IWRM Implementation database. UN SDG 6.5.1 database contains a category on financing which assesses different aspects related to budgeting and financing made available and used for water resources development and management from various sources.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
8.3 - Financing for Water Resource Development and Management (SDG 6.5.1) (rank)	19	This risk indicator is based on the average 'Financing' score of UN SDG 6.5.1. Degree of IWRM Implementation database. UN SDG 6.5.1 database contains a category on financing which assesses different aspects related to budgeting and financing made available and used for water resources development and management from various sources.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
9.1 - Cultural Diversity (score)	1.00	Water is a social and cultural good. The cultural diversity risk indicator was included in order to acknowledge that businesses face reputational risk due to the importance of freshwater for indigenous and traditional people in their daily life, religion and culture. This risk indicator is based on Oviedo and Larsen (2000) data set, which mapped the world's ethnolinguistic groups onto the WWF map of the world's ecoregions. This cross-mapping showed for the very first time the significant overlap that exists between the global geographic distribution of biodiversity and that of linguistic diversity.	Oviedo, G., Maffi, L., & Larsen, P. B. (2000). Indigenous and traditional peoples of the world and ecoregion conservation: An integrated approach to conserving the world's biological and cultural diversity. Gland: WWF (World Wide Fund for Nature) International.
9.1 - Cultural Diversity (rank)	129	Water is a social and cultural good. The cultural diversity risk indicator was included in order to acknowledge that businesses face reputational risk due to the importance of freshwater for indigenous and traditional people in their daily life, religion and culture. This risk indicator is based on Oviedo and Larsen (2000) data set, which mapped the world's ethnolinguistic groups onto the WWF map of the world's ecoregions. This cross-mapping showed for the very first time the significant overlap that exists between the global geographic distribution of biodiversity and that of linguistic diversity.	Oviedo, G., Maffi, L., & Larsen, P. B. (2000). Indigenous and traditional peoples of the world and ecoregion conservation: An integrated approach to conserving the world's biological and cultural diversity. Gland: WWF (World Wide Fund for Nature) International.
10.1 - Freshwater Endemism (score)	1.62	The underlying data set for this risk indicator comes from the Freshwater Ecoregions of the World (FEOW) 2015 data developed by WWF and TNC. Companies operating in basins with higher number of endemic fish species are exposed to higher reputational risks.	WWF & TNC (2015). Freshwater Ecoregions of the World.

Country Overview - Egypt

Indicator	Value	Description	Source
10.1 - Freshwater Endemism (rank)	184	The underlying data set for this risk indicator comes from the Freshwater Ecoregions of the World (FEOW) 2015 data developed by WWF and TNC. Companies operating in basins with higher number of endemic fish species are exposed to higher reputational risks.	WWF & TNC (2015). Freshwater Ecoregions of the World.
10.2 - Freshwater Biodiversity Richness (score)	2.09	The underlying data set for this risk indicator comes from the Freshwater Ecoregions of the World (FEOW) 2015 data developed by WWF and TNC. Count of fish species is used as a representation of freshwater biodiversity richness. Companies operating in basins with higher number of fish species are exposed to higher reputational risks.	WWF & TNC (2015). Freshwater Ecoregions of the World.
10.2 - Freshwater Biodiversity Richness (rank)	141	The underlying data set for this risk indicator comes from the Freshwater Ecoregions of the World (FEOW) 2015 data developed by WWF and TNC. Count of fish species is used as a representation of freshwater biodiversity richness. Companies operating in basins with higher number of fish species are exposed to higher reputational risks.	WWF & TNC (2015). Freshwater Ecoregions of the World.
11.1 - National Media Coverage (score)	4.00	This risk indicator is based on joint qualitative research by WWF and Tecnomia (Typsa Group). It indicates how aware local residents typically are of water-related issues due to national media coverage. The status of the river basin (e.g., scarcity and pollution) is taken into account, as well as the importance of water for livelihoods (e.g., food and shelter).	WWF & Tecnomia (TYPESA Group)
11.1 - National Media Coverage (rank)	18	This risk indicator is based on joint qualitative research by WWF and Tecnomia (Typsa Group). It indicates how aware local residents typically are of water-related issues due to national media coverage. The status of the river basin (e.g., scarcity and pollution) is taken into account, as well as the importance of water for livelihoods (e.g., food and shelter).	WWF & Tecnomia (TYPESA Group)
11.2 - Global Media Coverage (score)	4.00	This risk indicator is based on joint qualitative research by WWF and Tecnomia (Typsa Group). It indicates how aware people are of water-related issues due to global media coverage. Familiarity to and media coverage of the region and regional water-related disasters are taken into account.	WWF & Tecnomia (TYPESA Group)
11.2 - Global Media Coverage (rank)	11	This risk indicator is based on joint qualitative research by WWF and Tecnomia (Typsa Group). It indicates how aware people are of water-related issues due to global media coverage. Familiarity to and media coverage of the region and regional water-related disasters are taken into account.	WWF & Tecnomia (TYPESA Group)

Country Overview - Egypt

Indicator	Value	Description	Source
12.1 - Conflict News Events (RepRisk) (score)	4.00	This risk indicator is based on 2018 data collected by RepRisk on counts and registers of documented negative incidents, criticism and controversies that can affect a company's reputational risk. These negative news events are labelled per country and industry class.	RepRisk & WWF (2019). Due diligence database on ESG and business conduct risks. RepRisk.
12.1 - Conflict News Events (RepRisk) (rank)	13	This risk indicator is based on 2018 data collected by RepRisk on counts and registers of documented negative incidents, criticism and controversies that can affect a company's reputational risk. These negative news events are labelled per country and industry class.	RepRisk & WWF (2019). Due diligence database on ESG and business conduct risks. RepRisk.
12.2 - Hydro-political Risk (score)	3.06	This risk indicator is based on the assessment of hydro-political risk by Farinosi et al. (2018). More specifically, it is based on the results of spatial modelling by Farinosi et al. (2018) that determined the main parameters affecting water cross-border conflicts and calculated the likelihood of hydro-political issues.	Farinosi, F., Giupponi, C., Reynaud, A., Ceccherini, G., Carmona-Moreno, C., De Roo, A., ... & Bidoglio, G. (2018). An innovative approach to the assessment of hydro-political risk: A spatially explicit, data driven indicator of hydro-political issues. <i>Global environmental change</i> , 52, 286-313.
12.2 - Hydro-political Risk (rank)	26	This risk indicator is based on the assessment of hydro-political risk by Farinosi et al. (2018). More specifically, it is based on the results of spatial modelling by Farinosi et al. (2018) that determined the main parameters affecting water cross-border conflicts and calculated the likelihood of hydro-political issues.	Farinosi, F., Giupponi, C., Reynaud, A., Ceccherini, G., Carmona-Moreno, C., De Roo, A., ... & Bidoglio, G. (2018). An innovative approach to the assessment of hydro-political risk: A spatially explicit, data driven indicator of hydro-political issues. <i>Global environmental change</i> , 52, 286-313.
Population, total (#)	95688681	Population, total	The World Bank 2018, Data , homepage accessed 20/04/2018
GDP (current US\$)	332791045964	GDP (current US\$)	The World Bank 2018, Data , homepage accessed 20/04/2018
EPI 2018 score (0-100)	61.21	Environmental Performance Index	
WGI -Voice and Accountability (0-100)	0.00	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, <i>The Worldwide Governance Indicators: Methodology and Analytical Issues</i> (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132

Country Overview - Egypt

Indicator	Value	Description	Source
WGI -Political stability no violence (0-100)	0.00	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, The Worldwide Governance Indicators: Methodology and Analytical Issues (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132
WGI - Government Effectiveness (0-100)	0.00	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, The Worldwide Governance Indicators: Methodology and Analytical Issues (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132
WGI - Regulatory Quality (0-100)	0.00	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, The Worldwide Governance Indicators: Methodology and Analytical Issues (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132
WGI - Rule of Law (0-100)	0.00	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, The Worldwide Governance Indicators: Methodology and Analytical Issues (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132
WGI - Control of Corruption (0-100)	0.00	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, The Worldwide Governance Indicators: Methodology and Analytical Issues (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132

Country Overview - Egypt

Indicator	Value	Description	Source
WRI BWS all industries (0-5)	1.33	WRI Baseline Water Stress (BWS)	Gassert, F., P. Reig, T. Luo, and A. Maddocks. 2013. "Aqueduct country and river basin rankings: a weighted aggregation of spatially distinct hydrological indicators." Working paper. Washington, DC: World Resources Institute, December 2013. Available online at http://wri.org/publication/aqueduct-country-river-basin-rankings .
WRI BWS Ranking (1=very high)	102	WRI Baseline Water Stress (BWS)	Gassert, F., P. Reig, T. Luo, and A. Maddocks. 2013. "Aqueduct country and river basin rankings: a weighted aggregation of spatially distinct hydrological indicators." Working paper. Washington, DC: World Resources Institute, December 2013. Available online at http://wri.org/publication/aqueduct-country-river-basin-rankings .
Baseline Water Stress (BWS) - 2020 BAU (1=very high)	88	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .
Baseline Water Stress (BWS) - 2020 Optimistic (increasing rank describes lower risk)	89	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .
Baseline Water Stress (BWS) - 2020 Pessimistic (increasing rank describes lower risk)	88	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .

Country Overview - Egypt

Indicator	Value	Description	Source
Baseline Water Stress (BWS) - 2030 BAU (increasing rank describes lower risk)	89	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .
Baseline Water Stress (BWS) - 2030 Optimistic (increasing rank describes lower risk)	89	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .
Baseline Water Stress (BWS) - 2030 Pessimistic (increasing rank describes lower risk)	89	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .
Baseline Water Stress (BWS) - 2040 BAU (increasing rank describes lower risk)	89	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .
Baseline Water Stress (BWS) - 2040 Optimistic (increasing rank describes lower risk)	90	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .
Baseline Water Stress (BWS) - 2040 Pessimistic (increasing rank describes lower risk)	89	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .

Country Overview - Egypt

Indicator	Value	Description	Source
Total water footprint of national consumption (m ³ /a/cap)	1341.02	WFN Water Footprint Data	Mekonnen, M.M. and Hoekstra, A.Y. (2011) National water footprint accounts: The green, blue and grey water footprint of production and consumption, Value of Water Research Report Series No. 50, UNESCO-IHE, Delft, the Netherlands. http://www.waterfootprint.org/Reports/Report50-NationalWaterFootprints-Vol1.pdf
Ratio external / total water footprint (%)	28.51	WFN Water Footprint Data	Mekonnen, M.M. and Hoekstra, A.Y. (2011) National water footprint accounts: The green, blue and grey water footprint of production and consumption, Value of Water Research Report Series No. 50, UNESCO-IHE, Delft, the Netherlands. http://www.waterfootprint.org/Reports/Report50-NationalWaterFootprints-Vol1.pdf
Area equipped for full control irrigation: total (1000 ha)	3610.00	Aquastat - Irrigation	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Area equipped for irrigation: total (1000 ha)	3610.00	Aquastat - Irrigation	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
% of the area equipped for irrigation actually irrigated (%)	100.00	Aquastat - Irrigation	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Electricity production from hydroelectric sources (% of total)	8.14	World Development Indicators	The World Bank 2018, Data , homepage accessed 20/04/2018
Total internal renewable water resources (IRWR) (10 ⁹ m ³ /year)	1.80	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Total internal renewable water resources (IRWR) (10 ⁹ m ³ /year)	56.50	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Water resources: total external renewable (10 ⁹ m ³ /year)	1.80	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13

Country Overview - Egypt

Indicator	Value	Description	Source
Total renewable water resources (10 ⁹ m ³ /year)	58.30	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Dependency ratio (%)	96.91	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Total renewable water resources per capita (m ³ /inhab/year)	637.10	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
World happiness [0-8]	4.42	WorldHappinessReport.org	World Happiness Report, homepage accessed 20/04/2018

Country Aspects

1. PHYSICAL ASPECTS

1.1. WATER RESOURCES

1.1.1. WATER RESOURCES

The Egyptian territory comprises the following river basins:

- The Northern Interior Basin in the east and southeast of the country, covering 520,881km² or 52 per cent of the total area of the country. A sub-basin of the Northern Interior Basin is the Qattara Depression;
- The Nile Basin, covering 326,751km² (33 per cent) in the central part of the country in the form of a broad north-south strip;
- The Mediterranean Coast Basin, covering 65,568km² (6 per cent); and
- The Northeast Coast Basin, a narrow strip of 88,250km² along the coast of the Red Sea (8 per cent).

The River Nile is the main source of water for Egypt, with an annual allocated flow of 55.5km³/year under the Nile Waters Agreement of 1959. Internal renewable surface water resources are estimated at 0.5km³/year. This brings total actual renewable surface water resources to 56km³/year. Internal renewable groundwater resources are estimated at 1.3km³/year. The overlap between surface water and groundwater being considered negligible, the total actual renewable water resources of the country are thus 57.3km³/year. The Nubian Sandstone aquifer located under the Western Desert is considered an important groundwater source, but this is fossil groundwater. The main source of internal recharge is percolation from irrigation water in the Valley and the Delta.

All drainage water in Upper Egypt, south of Cairo, flows back into the Nile and the irrigation canals; this amount is estimated at 4km³/year. Drainage water in the Nile Delta is estimated at 14km³/year. Treated municipal wastewater in 2001/02 was estimated at 2.97km³/year. There are several desalination plants on the coasts of the Red Sea and the Mediterranean to provide water for seaside resorts and hotels; total production in 2002 was estimated at 100 million m³. Estimates of the potential of non-renewable groundwater in the eastern and western deserts, mainly from the Nubian Sandstone aquifer, vary from 3.8km³/year to 0.6km³/year; the latter estimate is defined as an indicator of exploitability over a period of time, where the time is not given.

1.1.2. WATER USE

Total water withdrawal in 2000 was estimated at 68.3km³. This included 59km³ for agriculture (86 per cent), 5.3km³ for municipalities (8 per cent) and 4km³ for industry (6 per cent). Apart from that, 4km³ were used for navigation and hydropower.

Groundwater extraction in 2000 was 7.043km³ comprising:

- 6.127km³ from the Nile Basin (seepage waters);
- 0.825km³ from the eastern and western deserts, i.e. mainly the Nubian Sandstone aquifer; and
- 0.091km³ from shallow wells in Sinai and on the northwestern coast.

Reuse of agricultural drainage water, returned to the rivers in irrigation amounted to 4.84km³/yr in 2001/02. Of the 2.97km³/yr of treated wastewater, 1.5km³/yr is reused for irrigation, while the rest is pumped into main drains where it mixes with drainage water and is then used for irrigation. Treated wastewater is usually used for landscape irrigation of trees in urban areas and along roads.

1.2. WATER QUALITY, ECOSYSTEMS AND HUMAN HEALTH

Major environmental problems are increasing soil salination below Aswan High Dam; desertification; oil pollution threatening coral reefs, beaches, and marine habitats; other water pollution from agricultural pesticides, raw sewage, and industrial effluents; the limited natural fresh water resources away from the Nile, which is the only perennial water source; rapid growth in population overstraining the Nile and natural resources.

Salinity and waterlogging are now under control in about 80 per cent of the affected areas following the installation of drainage systems. This has led to a reduction in saline areas from about 12,000km³ in 1972 to 2,500km³ at present. Seawater intrusion is a problem in the northern part of the Delta, where groundwater becomes brackish to saline. About half of the Delta contains brackish to saline groundwater, and no water is pumped in the northern Delta. Average salinity of groundwater in the Delta was between 680 and 1,170 mg/litre in 2000/01. It is said that the rice belt is a major defence factor to keep salinity under control in the northern areas of the Nile Delta, as well as to stop the invasion of seawater further inland in the Delta aquifer.

The salinity of agricultural drainage water is higher in winter, especially downstream, because less water is used for irrigation. In the new Al Salam Canal, drainage water is mixed with Nile water at a ratio of 1:1, and the salinity of the mixed water is within safe levels.

The main challenge for the sustainability of water resources is the control of water pollution. The Ministry of the Environment is observing the enforcement of the new legislation regarding the treatment of industrial and municipal wastewater. MALR is also advocating organic farming and limiting the use of chemical fertilizers and pesticides to reduce crop, soil and water pollution. In addition, present policy is to minimize the use of herbicides and to depend mainly on the mechanized control of submerged weeds and water hyacinths.

Bilharzia, or schistosomiasis, is still a common disease in rural areas in Egypt but its occurrence has greatly decreased with the provision of improved drinking water to most rural areas, periodic examination of school children, free medical treatment and extension programmes to educate people on ways of protecting against the disease. The Ministry of Health and Population

announced that bilharzia cases in the examined samples of rural population in 2001 were only about 4 per cent. Malaria is rare in Egypt.

2. GOVERNANCE ASPECTS

2.1. WATER INSTITUTIONS

The Ministry of Water Resources and Irrigation (MWRI) is in charge of water resources research, development and distribution, and undertakes the construction, operation and maintenance of the irrigation and drainage networks. Specifications and permits for groundwater well drilling are also the responsibility of MWRI. Within MWRI, the following sectors and departments are of importance:

- The Planning Sector is responsible at central level for data collection, processing and analysis for planning and monitoring investment projects;
 - The Sector of Public Works and Water Resources coordinates water resources development works;
 - The Nile Water Sector is in charge of cooperation with Sudan and other nilotic countries;
 - The Irrigation Department provides technical guidance and monitoring of irrigation development, including dams; and
 - The Mechanical and Electrical Department is in charge of the construction and maintenance of pumping stations for irrigation and drainage.
- Further to the above institutions, other public authorities are directly related to MWRI:
- The High Dam Authority is responsible for dam operation;
 - The Drainage Authority is responsible for the construction and maintenance of tile and open drains; and
 - The National Water Research Center comprises 12 institutes and is the scientific body of MWRI for all aspects related to water resources management.

2.2. WATER MANAGEMENT

The Government has indicated its intent to shift emphasis from its role as the central (or sole) actor in developing and managing water supply systems, towards promoting participatory approaches in which water users will play an active role in the management of irrigation systems and cost sharing. Important institutional and legislative measures have been taken recently to promote the establishment of sustainable participatory irrigation management (PIM) associations. However, despite these measures, the development of water users' associations (WUAs) as effective partners in irrigation management remains at an early stage. In the new lands, the concept of PIM is not yet effectively operational for a variety of economic, financial and institutional reasons.

While most settlers recognize the importance of WUAs in the equitable distribution of available water, uneven water availability, either due to design shortcomings or to lax enforcement of rules against excess abstraction by front-end water users, has acted as a disincentive to the successful

operation of WUAs in many instances.

2.3. WATER POLICY AND LEGAL FRAMEWORK

The main water and irrigation strategy is concerned with the development and conservation of water resources. This is done through adopting water rotation for irrigation canals, limiting the rice growing area, lining irrigation canals in sandy areas and prohibiting surface irrigation in the new developed areas outside the Nile basin.

Recent water resources policies include different structural and several non-structural measures. Structural measures include: irrigation structures rehabilitation; improvement of the irrigation system; installation of water level monitoring devices linked to the telemetry system; and expansion of the tile drainage system. Non-structural measures include: expansion of water user associations (WUAs) for irrigation ditches; establishment of water boards on branch canals; promotion of public awareness programmes; and involvement of stakeholders.

The legal basis for irrigation and drainage is set out in Law No. 12/1984 and its supplementary Law No. 213/1994 which define the use and management of public and private sector irrigation and drainage systems including main canals, feeders and drains. The laws also provide legal directions for the operation and maintenance of public and private waterways and specify arrangements for cost recovery in irrigation and drainage networks.

The most recent water policy was drafted in 1993. It included several strategies to ensure satisfying the demands of all water users and expanding the existing agricultural area at that time of 7.8 million feddan (about 31,200km³) by an additional 1.4 million feddan (about 5,600km³).

3. GEOPOLITICAL ASPECTS

Under the 1959 Nile Waters Agreement between Egypt and Sudan, Egypt's share of the Nile flow is 55.5km³/year. The agreement was based on the average flow of the Nile during the 1900-1959 period, which was 84km³/year at Aswan. Average annual evaporation and other losses from the Aswan High Dam and reservoir (Lake Nasser) were estimated at 10km³/year, leaving a net usable flow of 74km³/year, of which 18.5km³/year was allocated to Sudan and 55.5km³/year to Egypt.

In 1998, recognizing that cooperative development was the best way to bring mutual benefits to the region, all riparian countries (except Eritrea which had observer status only) joined in a dialogue to create a regional partnership to facilitate the common pursuit of sustainable development and management of Nile waters. The transitional mechanism, the Nile Basin Initiative (NBI), was officially launched in February 1999 in Dar es Salaam, in the United Republic of Tanzania, by the Council of Ministers of Water Affairs of the Nile Basin States. The shared vision of the NBI is "to achieve sustainable socio-economic development through the equitable utilization of and benefit from the common Nile Basin water resources".

The first meeting of the International Consortium for Cooperation on the Nile (ICCON) took place in 2001 in Geneva, Switzerland to celebrate cooperation between the ten countries of the Nile Basin and to establish partnerships leading to sustainable development and management of the Nile River for the benefit of all. The ICCON's first meeting was a major milestone for the NBI as it

Country Overview - Egypt

brought together, for the first time, ministers and senior officials from Nile Basin countries with a broad range of bilateral and multilateral donors and other interested parties from civil society, professional organizations, the media and NGOs.

If conditions permit the completion of the development projects on the Upper Nile, Egypt's share in the Nile water will increase by 9km³. This amount includes 1.9km³ and 1.6km³ respectively from the first and second phases of the Jonglei canal project in southern Sudan. Two other projects in the upstream swamps are expected to provide 5.5km³.

The vast Nubian Sandstone aquifer is shared with the Libyan Arab Jamahiriya, Sudan and Chad.