

Water Indicators

Indicator	Value	Description	Source
Overall Basin Risk (score)	2.03	Overall Basin Risk (score)	
Overall Basin Risk (rank)	181	Overall Basin Risk (rank)	
Physical risk (score)	2.02	Physical risk (score)	
Physical risk (rank)	167	Physical risk (rank)	
Regulatory risk (score)	1.64	Regulatory risk (score)	
Regulatory risk (rank)	167	Regulatory risk (rank)	
Reputation risk (score)	2.44	Reputation risk (score)	
Reputation risk (rank)	136	Reputation risk (rank)	
1. Quantity - Scarcity (score)	1.39	1. Quantity - Scarcity (score)	
1. Quantity - Scarcity (rank)	175	1. Quantity - Scarcity (rank)	
2. Quantity - Flooding (score)	1.92	2. Quantity - Flooding (score)	
2. Quantity - Flooding (rank)	166	2. Quantity - Flooding (rank)	
3. Quality (score)	3.26	3. Quality (score)	
3. Quality (rank)	75	3. Quality (rank)	
4. Ecosystem Service Status (score)	2.83	4. Ecosystem Service Status (score)	
4. Ecosystem Service Status (rank)	64	4. Ecosystem Service Status (rank)	
5. Enabling Environment (Policy & Laws) (score)	1.45	5. Enabling Environment (Policy & Laws) (score)	
5. Enabling Environment (Policy & Laws) (rank)	158	5. Enabling Environment (Policy & Laws) (rank)	
6. Institutions and Governance (score)	2.00	6. Institutions and Governance (score)	
6. Institutions and Governance (rank)	171	6. Institutions and Governance (rank)	
7. Management Instruments (score)	1.42	7. Management Instruments (score)	
7. Management Instruments (rank)	175	7. Management Instruments (rank)	
8 - Infrastructure & Finance (score)	1.65	8 - Infrastructure & Finance (score)	
8 - Infrastructure & Finance (rank)	122	8 - Infrastructure & Finance (rank)	
9. Cultural Diversity (score)	3.00	9. Cultural importance (score)	
9. Cultural Diversity (rank)	85	9. Cultural importance (rank)	
10. Biodiversity Importance (score)	2.64	10. Biodiversity importance (score)	

Country Overview - Latvia

Indicator	Value	Description	Source
10. Biodiversity Importance (rank)	157	10. Biodiversity importance (rank)	
11. Media Scrutiny (score)	2.10	11. Media Scrutiny (score)	
11. Media Scrutiny (rank)	179	11. Media Scrutiny (rank)	
12. Conflict (score)	2.55	12. Conflict (score)	
12. Conflict (rank)	85	12. Conflict (rank)	
1.0 - Aridity (score)	1.00	The aridity risk indicator is based on the Global Aridity Index (Global-Aridity) and Global Potential Evapo-Transpiration (Global-PET) Geospatial data sets by Trabucco and Zomer (2009). These data sets provide information about the potential availability of water in regions with low water demand, thus they are used in the Water Risk Filter 5.0 to better account for deserts and other arid areas in the risk assessment.	Trabucco, A., & Zomer, R. J. (2009). Global potential evapo-transpiration (Global-PET) and global aridity index (Global-Aridity) geodatabase. CGIAR consortium for spatial information.
1.0 - Aridity (rank)	184	The aridity risk indicator is based on the Global Aridity Index (Global-Aridity) and Global Potential Evapo-Transpiration (Global-PET) Geospatial data sets by Trabucco and Zomer (2009). These data sets provide information about the potential availability of water in regions with low water demand, thus they are used in the Water Risk Filter 5.0 to better account for deserts and other arid areas in the risk assessment.	Trabucco, A., & Zomer, R. J. (2009). Global potential evapo-transpiration (Global-PET) and global aridity index (Global-Aridity) geodatabase. CGIAR consortium for spatial information.
1.1 - Water Depletion (score)	1.24	The water depletion risk indicator is based on annual average monthly net water depletion from Brauman et al. (2016). Their analysis is based on model outputs from the newest version of the integrated water resources model WaterGAP3 which measures water depletion as the ratio of water consumption-to-availability.	Brauman, K. A., Richter, B. D., Postel, S., Malsy, M., & Flörke, M. (2016). Water depletion: An improved metric for incorporating seasonal and dry-year water scarcity into water risk assessments. <i>Elem Sci Anth</i> , 4.
1.1 - Water Depletion (rank)	121	The water depletion risk indicator is based on annual average monthly net water depletion from Brauman et al. (2016). Their analysis is based on model outputs from the newest version of the integrated water resources model WaterGAP3 which measures water depletion as the ratio of water consumption-to-availability.	Brauman, K. A., Richter, B. D., Postel, S., Malsy, M., & Flörke, M. (2016). Water depletion: An improved metric for incorporating seasonal and dry-year water scarcity into water risk assessments. <i>Elem Sci Anth</i> , 4.
1.2 - Baseline Water Stress (score)	1.12	World Resources Institute's Baseline Water Stress measures the ratio of total annual water withdrawals to total available annual renewable supply, accounting for upstream consumptive use. A higher percentage indicates more competition among users.	Hofste, R., Kuzma, S., Walker, S., ... & Sutanudjaja, E.H. (2019). <i>Aqueduct 3.0: Updated decision relevant global water risk indicators</i> . Technical note. Washington, DC: World Resources Institute.

Country Overview - Latvia

Indicator	Value	Description	Source
1.2 - Baseline Water Stress (rank)	142	World Resources Institute's Baseline Water Stress measures the ratio of total annual water withdrawals to total available annual renewable supply, accounting for upstream consumptive use. A higher percentage indicates more competition among users.	Hofste, R., Kuzma, S., Walker, S., ... & Sutanudjaja, E.H. (2019). Aqueduct 3.0: Updated decision relevant global water risk indicators. Technical note. Washington, DC: World Resources Institute.
1.3 - Blue Water Scarcity (score)	1.00	The blue water scarcity risk indicator is based on Mekonnen and Hoekstra (2016) global assessment of blue water scarcity on a monthly basis and at high spatial resolution (grid cells of 30 × 30 arc min resolution). Blue water scarcity is calculated as the ratio of the blue water footprint in a grid cell to the total blue water availability in the cell. The time period analyzed in this study ranges from 1996 to 2005.	Mekonnen, M. M., & Hoekstra, A. Y. (2016). Four billion people facing severe water scarcity. <i>Science advances</i> , 2(2), e1500323.
1.3 - Blue Water Scarcity (rank)	188	The blue water scarcity risk indicator is based on Mekonnen and Hoekstra (2016) global assessment of blue water scarcity on a monthly basis and at high spatial resolution (grid cells of 30 × 30 arc min resolution). Blue water scarcity is calculated as the ratio of the blue water footprint in a grid cell to the total blue water availability in the cell. The time period analyzed in this study ranges from 1996 to 2005.	Mekonnen, M. M., & Hoekstra, A. Y. (2016). Four billion people facing severe water scarcity. <i>Science advances</i> , 2(2), e1500323.
1.4 - Projected Change in Water Discharge (by ~2050) (score)	1.56	This risk indicator is based on multi-model simulation that applies both global climate and hydrological models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP). To estimate the change at 2°C of global warming above 1980-2010 levels, simulated annual water discharge was averaged over a 31-year period with 2°C mean warming. Results are expressed in terms of relative change (%) in probability between present day (1980-2010) conditions and 2°C scenarios by 2050.	Schewe, J., Heinke, J., Gerten, D., Haddeland, I., Arnell, N. W., Clark, D. B., ... & Gosling, S. N. (2014). Multimodel assessment of water scarcity under climate change. <i>Proceedings of the National Academy of Sciences</i> , 111(9), 3245-3250.
1.4 - Projected Change in Water Discharge (by ~2050) (rank)	129	This risk indicator is based on multi-model simulation that applies both global climate and hydrological models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP). To estimate the change at 2°C of global warming above 1980-2010 levels, simulated annual water discharge was averaged over a 31-year period with 2°C mean warming. Results are expressed in terms of relative change (%) in probability between present day (1980-2010) conditions and 2°C scenarios by 2050.	Schewe, J., Heinke, J., Gerten, D., Haddeland, I., Arnell, N. W., Clark, D. B., ... & Gosling, S. N. (2014). Multimodel assessment of water scarcity under climate change. <i>Proceedings of the National Academy of Sciences</i> , 111(9), 3245-3250.

Country Overview - Latvia

Indicator	Value	Description	Source
1.5 - Drought Frequency Probability (score)	2.04	This risk indicator is based on the Standardized Precipitation and Evaporation Index (SPEI). Vicente-Serrano et al. (2010) developed this multi-scalar drought index applying both precipitation and temperature data to detect, monitor and analyze different drought types and impacts in the context of global warming. The mathematical calculations used for SPEI are similar to the Standard Precipitation Index (SPI), but it has the advantage to include the role of evapotranspiration.	Vicente-Serrano, S. M., Beguería, S., & López-Moreno, J. I. (2010). A multiscale drought index sensitive to global warming: the standardized precipitation evapotranspiration index. <i>Journal of climate</i> , 23(7), 1696-1718.
1.5 - Drought Frequency Probability (rank)	124	This risk indicator is based on the Standardized Precipitation and Evaporation Index (SPEI). Vicente-Serrano et al. (2010) developed this multi-scalar drought index applying both precipitation and temperature data to detect, monitor and analyze different drought types and impacts in the context of global warming. The mathematical calculations used for SPEI are similar to the Standard Precipitation Index (SPI), but it has the advantage to include the role of evapotranspiration.	Vicente-Serrano, S. M., Beguería, S., & López-Moreno, J. I. (2010). A multiscale drought index sensitive to global warming: the standardized precipitation evapotranspiration index. <i>Journal of climate</i> , 23(7), 1696-1718.
1.6 - Projected Change in Drought Occurrence (by ~2050) (score)	2.64	This risk indicator is based on multi-model simulation that applies both global climate and drought models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP) . A drought threshold for pre-industrial conditions was calculated based on time-series averages. Results are expressed in terms of relative change (%) in probability between pre-industrial and 2°C scenarios.	Frieler, K., Lange, S., Piontek, F., Reyer, C. P., Schewe, J., Warszawski, L., ... & Geiger, T. (2017). Assessing the impacts of 1.5 C global warming-simulation protocol of the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP2b). Geoscientific Model Development.
1.6 - Projected Change in Drought Occurrence (by ~2050) (rank)	186	This risk indicator is based on multi-model simulation that applies both global climate and drought models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP) . A drought threshold for pre-industrial conditions was calculated based on time-series averages. Results are expressed in terms of relative change (%) in probability between pre-industrial and 2°C scenarios.	Frieler, K., Lange, S., Piontek, F., Reyer, C. P., Schewe, J., Warszawski, L., ... & Geiger, T. (2017). Assessing the impacts of 1.5 C global warming-simulation protocol of the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP2b). Geoscientific Model Development.
2.1 - Estimated Flood Occurrence (score)	1.96	This risk indicator is based on the recurrence of floods within the 34-year time frame period of 1985 to 2019. The occurrence of floods within a given location was estimated using data from Flood Observatory, University of Colorado. The Flood Observatory use data derived from a wide variety of news, governmental, instrumental, and remote sensing source.	Brakenridge, G. R. (2019). Global active archive of large flood events. Dartmouth Flood Observatory, University of Colorado.
2.1 - Estimated Flood Occurrence (rank)	163	This risk indicator is based on the recurrence of floods within the 34-year time frame period of 1985 to 2019. The occurrence of floods within a given location was estimated using data from Flood Observatory, University of Colorado. The Flood Observatory use data derived from a wide variety of news, governmental, instrumental, and remote sensing source.	Brakenridge, G. R. (2019). Global active archive of large flood events. Dartmouth Flood Observatory, University of Colorado.

Country Overview - Latvia

Indicator	Value	Description	Source
2.2 - Projected Change in Flood Occurrence (by ~2050) (score)	1.23	This risk indicator is based on multi-model simulation that applies both global climate and drought models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP). The magnitude of the flood event was defined based on 100-year return period for pre-industrial conditions. Results are expressed in terms of change (%) in probability between pre-industrial and 2°C scenarios.	Frieler, K., Lange, S., Piontek, F., Reyer, C. P., Schewe, J., Warszawski, L., ... & Geiger, T. (2017). Assessing the impacts of 1.5 C global warming-simulation protocol of the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP2b). Geoscientific Model Development.
2.2 - Projected Change in Flood Occurrence (by ~2050) (rank)	176	This risk indicator is based on multi-model simulation that applies both global climate and drought models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP). The magnitude of the flood event was defined based on 100-year return period for pre-industrial conditions. Results are expressed in terms of change (%) in probability between pre-industrial and 2°C scenarios.	Frieler, K., Lange, S., Piontek, F., Reyer, C. P., Schewe, J., Warszawski, L., ... & Geiger, T. (2017). Assessing the impacts of 1.5 C global warming-simulation protocol of the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP2b). Geoscientific Model Development.
3.1 - Surface Water Contamination Index (score)	3.26	<p>The underlying data for this risk indicator is based on a broad suite of pollutants with well-documented direct or indirect negative effects on water security for both humans and freshwater biodiversity, compiled by Vörösmarty et al. (2010). The negative effects are specific to individual pollutants, ranging from impacts mediated by eutrophication such as algal blooms and oxygen depletion (e.g., caused by phosphorus and organic loading) to direct toxic effects (e.g., caused by pesticides, mercury).</p> <p>The overall Surface Water Contamination Index is calculated based on a range of key pollutants with different weightings according to the level of their negative effects on water security for both humans and freshwater biodiversity: soil salinization (8%), nitrogen (12%) and phosphorus (P, 13%) loading, mercury deposition (5%), pesticide loading (10%), sediment loading (17%), organic loading (as Biological Oxygen Demand, BOD; 15%), potential acidification (9%), and thermal alteration (11%).</p>	Vörösmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P., ... & Davies, P. M. (2010). Global threats to human water security and river biodiversity. <i>Nature</i> , 467(7315), 555.

Indicator	Value	Description	Source
3.1 - Surface Water Contamination Index (rank)	75	<p>The underlying data for this risk indicator is based on a broad suite of pollutants with well-documented direct or indirect negative effects on water security for both humans and freshwater biodiversity, compiled by Vörösmarty et al. (2010). The negative effects are specific to individual pollutants, ranging from impacts mediated by eutrophication such as algal blooms and oxygen depletion (e.g., caused by phosphorus and organic loading) to direct toxic effects (e.g., caused by pesticides, mercury).</p> <p>The overall Surface Water Contamination Index is calculated based on a range of key pollutants with different weightings according to the level of their negative effects on water security for both humans and freshwater biodiversity: soil salinization (8%), nitrogen (12%) and phosphorus (P, 13%) loading, mercury deposition (5%), pesticide loading (10%), sediment loading (17%), organic loading (as Biological Oxygen Demand, BOD; 15%), potential acidification (9%), and thermal alteration (11%).</p>	Vörösmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P., ... & Davies, P. M. (2010). Global threats to human water security and river biodiversity. <i>Nature</i> , 467(7315), 555.
4.1 - Fragmentation Status of Rivers (score)	2.25	<p>This risk indicator is based on the data set by Grill et al. (2019) mapping the world's free-flowing rivers. Grill et al. (2019) compiled a geometric network of the global river system and associated attributes, such as hydro-geometric properties, as well as pressure indicators to calculate an integrated connectivity status index (CSI). While only rivers with high levels of connectivity in their entire length are classified as free-flowing, rivers of CSI < 95% are considered as fragmented at a certain degree.</p>	Grill, G., Lehner, B., Thieme, M., Geenen, B., Tickner, D., Antonelli, F., ... & Macedo, H. E. (2019). Mapping the world's free-flowing rivers. <i>Nature</i> , 569(7755), 215.
4.1 - Fragmentation Status of Rivers (rank)	101	<p>This risk indicator is based on the data set by Grill et al. (2019) mapping the world's free-flowing rivers. Grill et al. (2019) compiled a geometric network of the global river system and associated attributes, such as hydro-geometric properties, as well as pressure indicators to calculate an integrated connectivity status index (CSI). While only rivers with high levels of connectivity in their entire length are classified as free-flowing, rivers of CSI < 95% are considered as fragmented at a certain degree.</p>	Grill, G., Lehner, B., Thieme, M., Geenen, B., Tickner, D., Antonelli, F., ... & Macedo, H. E. (2019). Mapping the world's free-flowing rivers. <i>Nature</i> , 569(7755), 215.
4.2 - Catchment Ecosystem Services Degradation Level (tree cover loss) (score)	4.63	<p>For this risk indicator, tree cover loss was applied as a proxy to represent catchment ecosystem services degradation since forests play an important role in terms of water regulation, supply and pollution control. The forest cover data is based on Hansen et al.'s global Landsat data at a 30-meter spatial resolution to characterize forest cover and change. The authors defined trees as vegetation taller than 5 meters in height, and forest cover loss as a stand-replacement disturbance, or a change from a forest to non-forest state, during the period 2000 – 2018.</p>	Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A. A., Tyukavina, A., ... & Kommareddy, A. (2013). High-resolution global maps of 21st-century forest cover change. <i>science</i> , 342(6160), 850-853.

Country Overview - Latvia

Indicator	Value	Description	Source
4.2 - Catchment Ecosystem Services Degradation Level (tree cover loss) (rank)	5	<p>For this risk indicator, tree cover loss was applied as a proxy to represent catchment ecosystem services degradation since forests play an important role in terms of water regulation, supply and pollution control.</p> <p>The forest cover data is based on Hansen et al.'s global Landsat data at a 30-meter spatial resolution to characterize forest cover and change. The authors defined trees as vegetation taller than 5 meters in height, and forest cover loss as a stand-replacement disturbance, or a change from a forest to non-forest state, during the period 2000 – 2018.</p>	Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A. A., Tyukavina, A., ... & Kommareddy, A. (2013). High-resolution global maps of 21st-century forest cover change. <i>science</i> , 342(6160), 850-853.
4.3 - Projected Impacts on Freshwater Biodiversity (score)	2.00	The study by Tedesco et al. (2013) to project changes [% increase or decrease] in extinction rate by ~2090 of freshwater fish due to water availability loss from climate change is used as a proxy to estimate the projected impacts on freshwater biodiversity.	Tedesco, P. A., Oberdorff, T., Cornu, J. F., Beauchard, O., Brosse, S., Dürr, H. H., ... & Hugueny, B. (2013). A scenario for impacts of water availability loss due to climate change on riverine fish extinction rates. <i>Journal of Applied Ecology</i> , 50(5), 1105-1115.
4.3 - Projected Impacts on Freshwater Biodiversity (rank)	149	The study by Tedesco et al. (2013) to project changes [% increase or decrease] in extinction rate by ~2090 of freshwater fish due to water availability loss from climate change is used as a proxy to estimate the projected impacts on freshwater biodiversity.	Tedesco, P. A., Oberdorff, T., Cornu, J. F., Beauchard, O., Brosse, S., Dürr, H. H., ... & Hugueny, B. (2013). A scenario for impacts of water availability loss due to climate change on riverine fish extinction rates. <i>Journal of Applied Ecology</i> , 50(5), 1105-1115.
5.1 - Freshwater Policy Status (SDG 6.5.1) (score)	2.00	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "National Water Resources Policy" indicator, which corresponds to one of the three national level indicators under the Enabling Environment category.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
5.1 - Freshwater Policy Status (SDG 6.5.1) (rank)	154	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "National Water Resources Policy" indicator, which corresponds to one of the three national level indicators under the Enabling Environment category.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
5.2 - Freshwater Law Status (SDG 6.5.1) (score)	1.00	<p>This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "National Water Resources Law(s)" indicator, which corresponds to one of the three national level indicators under the Enabling Environment category.</p> <p>For SDG 6.5.1, enabling environment depicts the conditions that help to support the implementation of IWRM, which includes legal and strategic planning tools for IWRM.</p>	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.

Indicator	Value	Description	Source
5.2 - Freshwater Law Status (SDG 6.5.1) (rank)	187	<p>This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation “National Water Resources Law(s)” indicator, which corresponds to one of the three national level indicators under the Enabling Environment category.</p> <p>For SDG 6.5.1, enabling environment depicts the conditions that help to support the implementation of IWRM, which includes legal and strategic planning tools for IWRM.</p>	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
5.3 - Implementation Status of Water Management Plans (SDG 6.5.1) (score)	2.00	<p>This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation “National IWRM plans” indicator, which corresponds to one of the three national level indicators under the Enabling Environment category.</p> <p>For SDG 6.5.1, enabling environment depicts the conditions that help to support the implementation of IWRM, which includes legal and strategic planning tools for IWRM.</p>	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
5.3 - Implementation Status of Water Management Plans (SDG 6.5.1) (rank)	164	<p>This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation “National IWRM plans” indicator, which corresponds to one of the three national level indicators under the Enabling Environment category.</p> <p>For SDG 6.5.1, enabling environment depicts the conditions that help to support the implementation of IWRM, which includes legal and strategic planning tools for IWRM.</p>	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
6.1 - Corruption Perceptions Index (score)	3.00	<p>This risk Indicator is based on the latest Transparency International's data: the Corruption Perceptions Index 2018. This index aggregates data from a number of different sources that provide perceptions of business people and country experts on the level of corruption in the public sector.</p>	Transparency International (2019). Corruption Perceptions Index 2018. Berlin: Transparency International.
6.1 - Corruption Perceptions Index (rank)	156	<p>This risk Indicator is based on the latest Transparency International's data: the Corruption Perceptions Index 2018. This index aggregates data from a number of different sources that provide perceptions of business people and country experts on the level of corruption in the public sector.</p>	Transparency International (2019). Corruption Perceptions Index 2018. Berlin: Transparency International.
6.2 - Freedom in the World Index (score)	1.00	<p>This risk indicator is based on Freedom House (2019), an annual global report on political rights and civil liberties, composed of numerical ratings and descriptive texts for each country and a select group of territories. The 2019 edition involved more than 100 analysts and more than 30 advisers with global, regional, and issue-based expertise to covers developments in 195 countries and 14 territories from January 1, 2018, through December 31, 2018.</p>	Freedom House (2019). Freedom in the world 2019. Washington, DC: Freedom House.

Indicator	Value	Description	Source
6.2 - Freedom in the World Index (rank)	185	This risk indicator is based on Freedom House (2019), an annual global report on political rights and civil liberties, composed of numerical ratings and descriptive texts for each country and a select group of territories. The 2019 edition involved more than 100 analysts and more than 30 advisers with global, regional, and issue-based expertise to covers developments in 195 countries and 14 territories from January 1, 2018, through December 31, 2018.	Freedom House (2019). Freedom in the world 2019. Washington, DC: Freedom House.
6.3 - Business Participation in Water Management (SDG 6.5.1) (score)	1.00	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "Business Participation in Water Resources Development, Management and Use" indicator, which corresponds to one of the six national level indicators under the Institutions and Participation category.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
6.3 - Business Participation in Water Management (SDG 6.5.1) (rank)	188	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "Business Participation in Water Resources Development, Management and Use" indicator, which corresponds to one of the six national level indicators under the Institutions and Participation category.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
7.1 - Management Instruments for Water Management (SDG 6.5.1) (score)	1.00	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "Sustainable and efficient water use management" indicator, which corresponds to one of the five national level indicators under the Management Instruments category. For SDG 6.5.1, management instruments refer to the tools and activities that enable decision-makers and users to make rational and informed choices between alternative actions.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
7.1 - Management Instruments for Water Management (SDG 6.5.1) (rank)	188	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "Sustainable and efficient water use management" indicator, which corresponds to one of the five national level indicators under the Management Instruments category. For SDG 6.5.1, management instruments refer to the tools and activities that enable decision-makers and users to make rational and informed choices between alternative actions.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.

Country Overview - Latvia

Indicator	Value	Description	Source
7.2 - Groundwater Monitoring Data Availability and Management (score)	1.00	This risk indicator is based on the data set by UN IGRAC (2019) to determine the level of availability of groundwater monitoring data at country level as groundwater management decisions rely strongly on data availability. The level of groundwater monitoring data availability for groundwater management is determined according to a combination of three criteria developed by WWF and IGRAC: 1) Status of country groundwater monitoring programme, 2) groundwater data availability for NGOs and 3) Public access to processed groundwater monitoring data.	UN IGRAC (2019). Global Groundwater Monitoring Network GGMM Portal. UN International Groundwater Resources Assessment Centre (IGRAC).
7.2 - Groundwater Monitoring Data Availability and Management (rank)	185	This risk indicator is based on the data set by UN IGRAC (2019) to determine the level of availability of groundwater monitoring data at country level as groundwater management decisions rely strongly on data availability. The level of groundwater monitoring data availability for groundwater management is determined according to a combination of three criteria developed by WWF and IGRAC: 1) Status of country groundwater monitoring programme, 2) groundwater data availability for NGOs and 3) Public access to processed groundwater monitoring data.	UN IGRAC (2019). Global Groundwater Monitoring Network GGMM Portal. UN International Groundwater Resources Assessment Centre (IGRAC).
7.3 - Density of Runoff Monitoring Stations (score)	3.77	The density of monitoring stations for water quantity was applied as proxy to develop this risk indicator. The Global Runoff Data Base was used to estimate the number of monitoring stations per 1000km ² of the main river system (data base access date: May 2018).	BfG (2019). Global Runoff Data Base. German Federal Institute of Hydrology (BfG).
7.3 - Density of Runoff Monitoring Stations (rank)	64	The density of monitoring stations for water quantity was applied as proxy to develop this risk indicator. The Global Runoff Data Base was used to estimate the number of monitoring stations per 1000km ² of the main river system (data base access date: May 2018).	BfG (2019). Global Runoff Data Base. German Federal Institute of Hydrology (BfG).
8.1 - Access to Safe Drinking Water (score)	1.00	This risk indicator is based on the Joint Monitoring Programme for Water Supply, Sanitation and Hygiene (UNICEF/WHO) 2019 data. It provides estimates on the use of water, sanitation and hygiene by country for the period 2000-2017.	WHO & UNICEF (2019). Estimates on the use of water, sanitation and hygiene by country (2000-2017). Joint Monitoring Programme for Water Supply, Sanitation and Hygiene.
8.1 - Access to Safe Drinking Water (rank)	182	This risk indicator is based on the Joint Monitoring Programme for Water Supply, Sanitation and Hygiene (UNICEF/WHO) 2019 data. It provides estimates on the use of water, sanitation and hygiene by country for the period 2000-2017.	WHO & UNICEF (2019). Estimates on the use of water, sanitation and hygiene by country (2000-2017). Joint Monitoring Programme for Water Supply, Sanitation and Hygiene.
8.2 - Access to Sanitation (score)	2.00	This risk indicator is based on the Joint Monitoring Programme for Water Supply, Sanitation and Hygiene (UNICEF/WHO) 2019 data. It provides estimates on the use of water, sanitation and hygiene by country for the period 2000-2017.	WHO & UNICEF (2019). Estimates on the use of water, sanitation and hygiene by country (2000-2017). Joint Monitoring Programme for Water Supply, Sanitation and Hygiene.

Indicator	Value	Description	Source
8.2 - Access to Sanitation (rank)	114	This risk indicator is based on the Joint Monitoring Programme for Water Supply, Sanitation and Hygiene (UNICEF/WHO) 2019 data. It provides estimates on the use of water, sanitation and hygiene by country for the period 2000-2017.	WHO & UNICEF (2019). Estimates on the use of water, sanitation and hygiene by country (2000-2017). Joint Monitoring Programme for Water Supply, Sanitation and Hygiene.
8.3 - Financing for Water Resource Development and Management (SDG 6.5.1) (score)	3.00	This risk indicator is based on the average 'Financing' score of UN SDG 6.5.1. Degree of IWRM Implementation database. UN SDG 6.5.1 database contains a category on financing which assesses different aspects related to budgeting and financing made available and used for water resources development and management from various sources.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
8.3 - Financing for Water Resource Development and Management (SDG 6.5.1) (rank)	141	This risk indicator is based on the average 'Financing' score of UN SDG 6.5.1. Degree of IWRM Implementation database. UN SDG 6.5.1 database contains a category on financing which assesses different aspects related to budgeting and financing made available and used for water resources development and management from various sources.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
9.1 - Cultural Diversity (score)	3.00	Water is a social and cultural good. The cultural diversity risk indicator was included in order to acknowledge that businesses face reputational risk due to the importance of freshwater for indigenous and traditional people in their daily life, religion and culture. This risk indicator is based on Oviedo and Larsen (2000) data set, which mapped the world's ethnolinguistic groups onto the WWF map of the world's ecoregions. This cross-mapping showed for the very first time the significant overlap that exists between the global geographic distribution of biodiversity and that of linguistic diversity.	Oviedo, G., Maffi, L., & Larsen, P. B. (2000). Indigenous and traditional peoples of the world and ecoregion conservation: An integrated approach to conserving the world's biological and cultural diversity. Gland: WWF (World Wide Fund for Nature) International.
9.1 - Cultural Diversity (rank)	85	Water is a social and cultural good. The cultural diversity risk indicator was included in order to acknowledge that businesses face reputational risk due to the importance of freshwater for indigenous and traditional people in their daily life, religion and culture. This risk indicator is based on Oviedo and Larsen (2000) data set, which mapped the world's ethnolinguistic groups onto the WWF map of the world's ecoregions. This cross-mapping showed for the very first time the significant overlap that exists between the global geographic distribution of biodiversity and that of linguistic diversity.	Oviedo, G., Maffi, L., & Larsen, P. B. (2000). Indigenous and traditional peoples of the world and ecoregion conservation: An integrated approach to conserving the world's biological and cultural diversity. Gland: WWF (World Wide Fund for Nature) International.
10.1 - Freshwater Endemism (score)	2.21	The underlying data set for this risk indicator comes from the Freshwater Ecoregions of the World (FEOW) 2015 data developed by WWF and TNC. Companies operating in basins with higher number of endemic fish species are exposed to higher reputational risks.	WWF & TNC (2015). Freshwater Ecoregions of the World.

Country Overview - Latvia

Indicator	Value	Description	Source
10.1 - Freshwater Endemism (rank)	166	The underlying data set for this risk indicator comes from the Freshwater Ecoregions of the World (FEOW) 2015 data developed by WWF and TNC. Companies operating in basins with higher number of endemic fish species are exposed to higher reputational risks.	WWF & TNC (2015). Freshwater Ecoregions of the World.
10.2 - Freshwater Biodiversity Richness (score)	3.07	The underlying data set for this risk indicator comes from the Freshwater Ecoregions of the World (FEOW) 2015 data developed by WWF and TNC. Count of fish species is used as a representation of freshwater biodiversity richness. Companies operating in basins with higher number of fish species are exposed to higher reputational risks.	WWF & TNC (2015). Freshwater Ecoregions of the World.
10.2 - Freshwater Biodiversity Richness (rank)	97	The underlying data set for this risk indicator comes from the Freshwater Ecoregions of the World (FEOW) 2015 data developed by WWF and TNC. Count of fish species is used as a representation of freshwater biodiversity richness. Companies operating in basins with higher number of fish species are exposed to higher reputational risks.	WWF & TNC (2015). Freshwater Ecoregions of the World.
11.1 - National Media Coverage (score)	3.00	This risk indicator is based on joint qualitative research by WWF and Tecnomia (Tyspa Group). It indicates how aware local residents typically are of water-related issues due to national media coverage. The status of the river basin (e.g., scarcity and pollution) is taken into account, as well as the importance of water for livelihoods (e.g., food and shelter).	WWF & Tecnomia (TYPESA Group)
11.1 - National Media Coverage (rank)	178	This risk indicator is based on joint qualitative research by WWF and Tecnomia (Tyspa Group). It indicates how aware local residents typically are of water-related issues due to national media coverage. The status of the river basin (e.g., scarcity and pollution) is taken into account, as well as the importance of water for livelihoods (e.g., food and shelter).	WWF & Tecnomia (TYPESA Group)
11.2 - Global Media Coverage (score)	1.00	This risk indicator is based on joint qualitative research by WWF and Tecnomia (Tyspa Group). It indicates how aware people are of water-related issues due to global media coverage. Familiarity to and media coverage of the region and regional water-related disasters are taken into account.	WWF & Tecnomia (TYPESA Group)
11.2 - Global Media Coverage (rank)	186	This risk indicator is based on joint qualitative research by WWF and Tecnomia (Tyspa Group). It indicates how aware people are of water-related issues due to global media coverage. Familiarity to and media coverage of the region and regional water-related disasters are taken into account.	WWF & Tecnomia (TYPESA Group)

Country Overview - Latvia

Indicator	Value	Description	Source
12.1 - Conflict News Events (RepRisk) (score)	3.00	This risk indicator is based on 2018 data collected by RepRisk on counts and registers of documented negative incidents, criticism and controversies that can affect a company's reputational risk. These negative news events are labelled per country and industry class.	RepRisk & WWF (2019). Due diligence database on ESG and business conduct risks. RepRisk.
12.1 - Conflict News Events (RepRisk) (rank)	107	This risk indicator is based on 2018 data collected by RepRisk on counts and registers of documented negative incidents, criticism and controversies that can affect a company's reputational risk. These negative news events are labelled per country and industry class.	RepRisk & WWF (2019). Due diligence database on ESG and business conduct risks. RepRisk.
12.2 - Hydro-political Risk (score)	2.10	This risk indicator is based on the assessment of hydro-political risk by Farinosi et al. (2018). More specifically, it is based on the results of spatial modelling by Farinosi et al. (2018) that determined the main parameters affecting water cross-border conflicts and calculated the likelihood of hydro-political issues.	Farinosi, F., Giupponi, C., Reynaud, A., Ceccherini, G., Carmona-Moreno, C., De Roo, A., ... & Bidoglio, G. (2018). An innovative approach to the assessment of hydro-political risk: A spatially explicit, data driven indicator of hydro-political issues. <i>Global environmental change</i> , 52, 286-313.
12.2 - Hydro-political Risk (rank)	115	This risk indicator is based on the assessment of hydro-political risk by Farinosi et al. (2018). More specifically, it is based on the results of spatial modelling by Farinosi et al. (2018) that determined the main parameters affecting water cross-border conflicts and calculated the likelihood of hydro-political issues.	Farinosi, F., Giupponi, C., Reynaud, A., Ceccherini, G., Carmona-Moreno, C., De Roo, A., ... & Bidoglio, G. (2018). An innovative approach to the assessment of hydro-political risk: A spatially explicit, data driven indicator of hydro-political issues. <i>Global environmental change</i> , 52, 286-313.
Population, total (#)	1960424	Population, total	The World Bank 2018, Data , homepage accessed 20/04/2018
GDP (current US\$)	27572698482	GDP (current US\$)	The World Bank 2018, Data , homepage accessed 20/04/2018
EPI 2018 score (0-100)	66.12	Environmental Performance Index	
WGI -Voice and Accountability (0-100)	59.52	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, <i>The Worldwide Governance Indicators: Methodology and Analytical Issues</i> (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132

Country Overview - Latvia

Indicator	Value	Description	Source
WGI -Political stability no violence (0-100)	74.38	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, The Worldwide Governance Indicators: Methodology and Analytical Issues (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132
WGI - Government Effectiveness (0-100)	78.85	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, The Worldwide Governance Indicators: Methodology and Analytical Issues (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132
WGI - Regulatory Quality (0-100)	83.65	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, The Worldwide Governance Indicators: Methodology and Analytical Issues (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132
WGI - Rule of Law (0-100)	80.29	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, The Worldwide Governance Indicators: Methodology and Analytical Issues (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132
WGI - Control of Corruption (0-100)	67.31	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, The Worldwide Governance Indicators: Methodology and Analytical Issues (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132

Country Overview - Latvia

Indicator	Value	Description	Source
WRI BWS all industries (0-5)	0.35	WRI Baseline Water Stress (BWS)	Gassert, F., P. Reig, T. Luo, and A. Maddocks. 2013. "Aqueduct country and river basin rankings: a weighted aggregation of spatially distinct hydrological indicators." Working paper. Washington, DC: World Resources Institute, December 2013. Available online at http://wri.org/publication/aqueduct-country-river-basin-rankings .
WRI BWS Ranking (1=very high)	144	WRI Baseline Water Stress (BWS)	Gassert, F., P. Reig, T. Luo, and A. Maddocks. 2013. "Aqueduct country and river basin rankings: a weighted aggregation of spatially distinct hydrological indicators." Working paper. Washington, DC: World Resources Institute, December 2013. Available online at http://wri.org/publication/aqueduct-country-river-basin-rankings .
Baseline Water Stress (BWS) - 2020 BAU (1=very high)	108	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .
Baseline Water Stress (BWS) - 2020 Optimistic (increasing rank describes lower risk)	109	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .
Baseline Water Stress (BWS) - 2020 Pessimistic (increasing rank describes lower risk)	108	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .

Country Overview - Latvia

Indicator	Value	Description	Source
Baseline Water Stress (BWS) - 2030 BAU (increasing rank describes lower risk)	107	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .
Baseline Water Stress (BWS) - 2030 Optimistic (increasing rank describes lower risk)	111	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .
Baseline Water Stress (BWS) - 2030 Pessimistic (increasing rank describes lower risk)	107	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .
Baseline Water Stress (BWS) - 2040 BAU (increasing rank describes lower risk)	109	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .
Baseline Water Stress (BWS) - 2040 Optimistic (increasing rank describes lower risk)	111	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .
Baseline Water Stress (BWS) - 2040 Pessimistic (increasing rank describes lower risk)	109	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct-projected-water-stress-country-rankings .

Country Overview - Latvia

Indicator	Value	Description	Source
Total water footprint of national consumption (m ³ /a/cap)	1796.76	WFN Water Footprint Data	Mekonnen, M.M. and Hoekstra, A.Y. (2011) National water footprint accounts: The green, blue and grey water footprint of production and consumption, Value of Water Research Report Series No. 50, UNESCO-IHE, Delft, the Netherlands. http://www.waterfootprint.org/Reports/Report50-NationalWaterFootprints-Vol1.pdf
Ratio external / total water footprint (%)	33.47	WFN Water Footprint Data	Mekonnen, M.M. and Hoekstra, A.Y. (2011) National water footprint accounts: The green, blue and grey water footprint of production and consumption, Value of Water Research Report Series No. 50, UNESCO-IHE, Delft, the Netherlands. http://www.waterfootprint.org/Reports/Report50-NationalWaterFootprints-Vol1.pdf
Area equipped for full control irrigation: total (1000 ha)	0.63	Aquastat - Irrigation	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Area equipped for irrigation: total (1000 ha)	0.63	Aquastat - Irrigation	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
% of the area equipped for irrigation actually irrigated (%)	65.08	Aquastat - Irrigation	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Electricity production from hydroelectric sources (% of total)	38.79	World Development Indicators	The World Bank 2018, Data , homepage accessed 20/04/2018
Total internal renewable water resources (IRWR) (10 ⁹ m ³ /year)	16.94	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Total internal renewable water resources (IRWR) (10 ⁹ m ³ /year)	18.00	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Water resources: total external renewable (10 ⁹ m ³ /year)	16.94	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13

Country Overview - Latvia

Indicator	Value	Description	Source
Total renewable water resources (10 ⁹ m ³ /year)	34.94	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Dependency ratio (%)	51.52	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Total renewable water resources per capita (m ³ /inhab/year)	17736.00	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
World happiness [0-8]	5.93	WorldHappinessReport.org	World Happiness Report, homepage accessed 20/04/2018

Country Aspects

1. PHYSICAL ASPECTS

1.1. WATER RESOURCES

1.1.1. WATER RESOURCES

Depending on the physical and geographical conditions, a large part of the river discharge comes from either snow melt, groundwater or direct surface runoff. About 50-55 per cent of the waters of the Daugava, Venta, Lielupe and Musa rivers is melted snow, while for the Gauja and Amata rivers it is 35-40 per cent. About 10-20 per cent of the flow of some tributaries of the Lielupe (Memele and Svete) and the Aiviekste tributary of the Daugava is fed by groundwater, while for the Daugava and Gauja rivers it is 35-40 per cent. In the Kurzeme peninsula and in the middle uplands, direct surface runoff accounts for 40 per cent of the flow of the rivers, while in the Zemgales plain it represents 20-30 per cent.

The country can be divided into eight river basins:

-The Daugava basin. Its total area is 87,900km², of which 28 per cent is located in Latvia. The Daugava River rises in the Russian Federation, flows through Belarus (where it is called the Western Dvina), enters Latvia in the southeast and flows northwest to the Gulf of Riga. Several tributaries enter the Daugava River inside Latvian territory, including four large ones: Ogre, Aiviekste, Dubna and Rezekne.

-The Gauja basin. Its total area is 8,900km², of which 88 per cent is situated in Latvia. The Gauja River rises in the Vidzeme upland and flows east, then turns northwest, becomes the border between Latvia and Estonia for a short distance, and then flows southwest to the Gulf of Riga.

-The Salaca basin. It covers the north of the country, near the border with Estonia. Its total area is 3,600km², of which 92 per cent is located in Latvia. The Salaca River rises in Lake Burtnieks in the north and flows west to the Gulf of Riga.

-The Lielupe basin. Its total area is 17,600km², of which 50 per cent is situated in Latvia. The Lielupe River rises in Lithuania, enters Latvia in the south and flows north to the Gulf of Riga through the most fertile regions of the country. It has many tributaries, the most important being the Memela, Jecava and Svete.

-The Venta basin. Its total area is 11,800km², of which 67 per cent is situated in Latvia. The Venta River rises in Lithuania, enters Latvia in the southwest and flows north through the Kurzeme lowland to the Baltic Sea. The Venta has many tributaries, but only one of them, the Abava River, exceeds 100km in length.

-The coastal basins between Lithuania and the Venta. Their total area is 5,100km². This area includes rivers such as the Barta, Durba, Riva and Uzava, which flow to the Baltic Sea.

-The basins within the coastal lowland, on the opposite shores of the Gulf of Riga. Their combined

area is 3,800km². This area includes rivers such as the Irbe, Stonde, Roja, Svetupe and Vitupe.

-The Velika basin. This basin consists of a number of smaller rivers flowing into the Velika in the Russian Federation. Its area within Latvia is 3,200km². The total discharge of the Velika amounts to 4.2km³/year, of which 16 per cent is generated within Latvia.

The total water resources are estimated at 16.54km³/year, incoming surface water resources at 18.709 km³/year.

The internal renewable groundwater resources are estimated at 2.2km³/year. Part of the groundwater flows to the sea or is withdrawn by wells, and part is drained by the surface network. That part of the groundwater flow which does not contribute to the total IRWR (overlap) is estimated at 2km³/year. Groundwater use is estimated at about 800,000m³/day. In some regions, rapid depletion of the water table is observed. Quite a large quantity is used by cities. In the Jūrmala area, close to the capital Riga, the groundwater is famed for its medicinal qualities

1.1.2. WATER USE

In 1994, the total water withdrawal for agricultural, domestic and industrial purposes was estimated at 285.2 million m³, of which about 12.7 per cent for agriculture. Other water use, including water use for hydropower, was 149.1 million m³.

In 1994, the total quantity of produced wastewater was 215.8 million m³, of which 119.4 million m³ were classed as clean without treatment, while 66.7 million m³ were treated to meet quality standards. The remaining 29.7 million m³ were not treated. The largest quantity of untreated wastewater flows into the Daugava and Lielupe river basins. In 1994, the total quantity of re-used treated wastewater amounted to 11.8 million m³.

1.2. WATER QUALITY, ECOSYSTEMS AND HUMAN HEALTH

Latvia's environment has benefited from a shift to service industries after the country regained independence. The main environmental priorities are the improvement of drinking water quality and the sewage system; household, and hazardous waste management; and reduction of air pollution. In 2001, Latvia closed the EU accession negotiation chapter on the environment by committing to full enforcement of EU environmental directives by 2010.

Eutrophication of water bodies and degradation of water ecosystems are priority environmental problems in Latvia, in relation to both inland and marine waters.

2. GOVERNANCE ASPECTS

2.1. WATER INSTITUTIONS

The following institutions are involved in water resources management:

-The Ministry of Agriculture is responsible for water management. It includes the Department of

Land Reclamation in Riga and local departments in every district. On behalf of the local departments, a land improvement specialist supervises the drainage networks of two or three communes. The funds allocated for the reconstruction and maintenance of the networks are managed by district departments and are allotted after consultations with farmers' organizations (mainly the Latvian Farmers Federation).

-The Ministry of Environmental Protection and Regional Development, with the Environmental Protection Department, collects information on the quantity and quality of water, and is responsible for preventing pollution of water, agricultural land and air.

-The Ministry of Power Engineering is responsible for hydroelectric power stations.

-The Latvian Hydro-meteorological Agency deals with recording the water quantities of the rivers, lakes and reservoirs.

-The Melioprojects Company VU in Riga keeps all the records concerning the process of land improvement. There are detailed maps (to a scale of 1:50 000 and 1:100 000) of particular districts with newly drained areas.

-The Latvian University of Agriculture in Jelgava carries out large-scale scientific research on irrigation and land reclamation.

-The Latvian State Research Institute, Agriculture Polymers and Water Management in Jelgava deals with matters of land reclamation, protection of the water against natural and anthropogenic pollution, and hydraulic modelling.

-The University of Latvia in Riga conducts research on hydrology, climate, the geography of water resources (Faculty of Geography) and in environmental sciences (Faculty of Geography, Centre for Environmental Studies). It collects statistics and information on water management, agriculture and irrigation.

2.2. WATER MANAGEMENT

A top priority for the 1995 National Environment Policy Plan for Latvia is the environmental protection of water courses in the Baltic Sea basin, since 85 per cent of the surface water resources are classed as either slightly polluted or polluted. Eutrophication is the most important problem.

Another task is to extend the agricultural period through faster draining during spring. Since 1991, there has been no drainage development at all in Latvia. Over 66,000km of drainage ditches (almost all the existing ditches) require rehabilitation. During the next ten years, it is planned to drain an additional 1,500 - 2,000km² in order to achieve self-sufficiency in cereals.

A priority of the EU Water Framework Directive (2000/60/EC) is integrated management of river drainage basins. This approach has been introduced in Latvia for water protection and management, and is one of the instruments in abatement of the impact of eutrophication.

2.3. WATER POLICY AND LEGAL FRAMEWORK

In the process of harmonization and implementation of EU legislation, the Latvian government has set 1 January 2003 as the date when Latvia can accept the role of an EU member.

In October 2000, the Cabinet of Ministers approved the negotiation position paper of Latvia in the chapter Environmental Protection. Initially, Latvia requested transition periods for 23 EU directives. After additional discussions and consultations with the European Commission, the number of requested transition periods was decreased to eight.

In order to implement the requirements of the Nitrates Directive (91/676/EEC), the Cabinet of Ministers Regulations On the Protection of Water and Soil Against Pollution Caused by Nitrates from Agricultural Sources was adopted.

The Dangerous Substances Directive (76/464/EEC) (and its daughter directives), the Groundwater Directive (80/68/EEC) and the Urban Wastewater Directive (91/271/EEC) have been implemented by The Cabinet of Ministers Regulations On Water Emissions of Pollutants. Full transposition of these Directives has been achieved.

The Directive on the Assessment of the Effects of Certain Public and Private Projects on the Environment (85/337/EEC and 97/11/EC) was implemented by the Law On Environmental Impact Assessment in October 1998. Amendments to the Law, dealing with outstanding requirements in respect of transboundary issues relating to environmental impact assessment, were adopted in May 2001 completing the implementation of the Directive.

The provisions of the Directive On the Freedom of Access to Information on the Environment (90/313/EEC) have fully been adopted in Latvian law through amendments to the Law On Environmental Protection, which were adopted by the parliament on 20 June 2000.

3. GEOPOLITICAL ASPECTS

Under Soviet administration, no agreements existed with neighbouring republics. Now that Latvia is independent, agreements are recognized as a necessity. An agreement with the Russian Federation and Belarus concerning the prevention of pollution of the water courses in the Daugava basin is under preparation, as is an agreement with Lithuania on the prevention of water pollution in the Lielupe, Venta and Barta rivers. No agreements on water sharing exist.

Only 44 per cent of the annual water discharge from rivers is contributed from the territory of Latvia. The other 56 per cent originates from Lithuania, Belarus and Russia, bringing with it pollution from other countries.

Transboundary water pollution is an important problem for internal waters, which is emphasized in the National Environmental Protection Policy Plan. The pollution originating from other countries contains significant levels of all pollutant groups, particularly nutrients and persistent substances.

The largest rivers receiving pollution from neighbouring countries are the Daugava, the Lielupe (Mēmele and Mūsa near the border), and the Venta.