

Water Indicators

Indicator	Value	Description	Source
Overall Basin Risk (score)	2.36	Overall Basin Risk (score)	
Overall Basin Risk (rank)	155	Overall Basin Risk (rank)	
Physical risk (score)	2.68	Physical risk (score)	
Physical risk (rank)	91	Physical risk (rank)	
Regulatory risk (score)	1.07	Regulatory risk (score)	
Regulatory risk (rank)	193	Regulatory risk (rank)	
Reputation risk (score)	2.70	Reputation risk (score)	
Reputation risk (rank)	89	Reputation risk (rank)	
1. Quantity - Scarcity (score)	1.76	1. Quantity - Scarcity (score)	
1. Quantity - Scarcity (rank)	140	1. Quantity - Scarcity (rank)	
2. Quantity - Flooding (score)	3.03	2. Quantity - Flooding (score)	
2. Quantity - Flooding (rank)	104	2. Quantity - Flooding (rank)	
3. Quality (score)	4.00	3. Quality (score)	
3. Quality (rank)	28	3. Quality (rank)	
4. Ecosystem Service Status (score)	3.79	4. Ecosystem Service Status (score)	
4. Ecosystem Service Status (rank)	4	4. Ecosystem Service Status (rank)	
5. Enabling Environment (Policy & Laws) (score)	1.10	5. Enabling Environment (Policy & Laws) (score)	
5. Enabling Environment (Policy & Laws) (rank)	170	5. Enabling Environment (Policy & Laws) (rank)	
6. Institutions and Governance (score)	1.00	6. Institutions and Governance (score)	
6. Institutions and Governance (rank)	189	6. Institutions and Governance (rank)	
7. Management Instruments (score)	1.15	7. Management Instruments (score)	
7. Management Instruments (rank)	192	7. Management Instruments (rank)	
8 - Infrastructure & Finance (score)	1.00	8 - Infrastructure & Finance (score)	
8 - Infrastructure & Finance (rank)	186	8 - Infrastructure & Finance (rank)	
9. Cultural Diversity (score)	3.00	9. Cultural importance (score)	
9. Cultural Diversity (rank)	75	9. Cultural importance (rank)	
10. Biodiversity Importance (score)	4.50	10. Biodiversity importance (score)	



Indicator	Value	Description	Source
10. Biodiversity Importance (rank)	23	10. Biodiversity importance (rank)	
11. Media Scrutiny (score)	2.55	11. Media Scrutiny (score)	
11. Media Scrutiny (rank)	139	11. Media Scrutiny (rank)	
12. Conflict (score)	2.09	12. Conflict (score)	
12. Conflict (rank)	137	12. Conflict (rank)	
1.0 - Aridity (score)	1.00	The aridity risk indicator is based on the Global Aridity Index (Global- Aridity) and Global Potential Evapo-Transpiration (Global-PET) Geospatial data sets by Trabucco and Zomer (2009). These data sets provide information about the potential availability of water in regions with low water demand, thus they are used in the Water Risk Filter 5.0 to better account for deserts and other arid areas in the risk assessment.	Trabucco, A., & Zomer, R. J. (2009). Global potential evapo-transpiration (Global-PET) and global aridity index (Global-Aridity) geo- database. CGIAR consortium for spatial information.
1.0 - Aridity (rank)	161	The aridity risk indicator is based on the Global Aridity Index (Global- Aridity) and Global Potential Evapo-Transpiration (Global-PET) Geospatial data sets by Trabucco and Zomer (2009). These data sets provide information about the potential availability of water in regions with low water demand, thus they are used in the Water Risk Filter 5.0 to better account for deserts and other arid areas in the risk assessment.	Trabucco, A., & Zomer, R. J. (2009). Global potential evapo-transpiration (Global-PET) and global aridity index (Global-Aridity) geo- database. CGIAR consortium for spatial information.
1.1 - Water Depletion (score)	1.00	The water depletion risk indicator is based on annual average monthly net water depletion from Brauman et al. (2016). Their analysis is based on model outputs from the newest version of the integrated water resources model WaterGAP3 which measures water depletion as the ratio of water consumption-to-availability.	Brauman, K. A., Richter, B. D., Postel, S., Malsy, M., & Flörke, M. (2016). Water depletion: An improved metric for incorporating seasonal and dry-year water scarcity into water risk assessments. Elem Sci Anth, 4.
1.1 - Water Depletion (rank)	183	The water depletion risk indicator is based on annual average monthly net water depletion from Brauman et al. (2016). Their analysis is based on model outputs from the newest version of the integrated water resources model WaterGAP3 which measures water depletion as the ratio of water consumption-to-availability.	Brauman, K. A., Richter, B. D., Postel, S., Malsy, M., & Flörke, M. (2016). Water depletion: An improved metric for incorporating seasonal and dry-year water scarcity into water risk assessments. Elem Sci Anth, 4.
1.2 - Baseline Water Stress (score)	3.03	World Resources Institute's Baseline Water Stress measures the ratio of total annual water withdrawals to total available annual renewable supply, accounting for upstream consumptive use. A higher percentage indicates more competition among users.	Hofste, R., Kuzma, S., Walker, S., & Sutanudjaja, E.H. (2019). Aqueduct 3.0: Updated decision relevant global water risk indicators. Technical note. Washington, DC: World Resources Institute.



Indicator	Value	Description	Source
1.2 - Baseline Water Stress (rank)	53	World Resources Institute's Baseline Water Stress measures the ratio of total annual water withdrawals to total available annual renewable supply, accounting for upstream consumptive use. A higher percentage indicates more competition among users.	Hofste, R., Kuzma, S., Walker, S., & Sutanudjaja, E.H. (2019). Aqueduct 3.0: Updated decision relevant global water risk indicators. Technical note. Washington, DC: World Resources Institute.
1.3 - Blue Water Scarcity (score)	1.00	The blue water scarcity risk indicator is based on Mekonnen and Hoekstra (2016) global assessment of blue water scarcity on a monthly basis and at high spatial resolution (grid cells of 30 × 30 arc min resolution). Blue water scarcity is calculated as the ratio of the blue water footprint in a grid cell to the total blue water availability in the cell. The time period analyzed in this study ranges from 1996 to 2005.	Mekonnen, M. M., & Hoekstra, A. Y. (2016). Four billion people facing severe water scarcity. Science advances, 2(2), e1500323.
1.3 - Blue Water Scarcity (rank)	181	The blue water scarcity risk indicator is based on Mekonnen and Hoekstra (2016) global assessment of blue water scarcity on a monthly basis and at high spatial resolution (grid cells of 30 × 30 arc min resolution). Blue water scarcity is calculated as the ratio of the blue water footprint in a grid cell to the total blue water availability in the cell. The time period analyzed in this study ranges from 1996 to 2005.	Mekonnen, M. M., & Hoekstra, A. Y. (2016). Four billion people facing severe water scarcity. Science advances, 2(2), e1500323.
1.4 - Projected Change in Water Discharge (by ~2050) (score)	2.00	This risk indicator is based on multi-model simulation that applies both global climate and hydrological models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP). To estimate the change at 2°C of global warming above 1980-2010 levels, simulated annual water discharge was averaged over a 31-year period with 2°C mean warming. Results are expressed in terms of relative change (%) in probability between present day (1980-2010) conditions and 2°C scenarios by 2050.	Schewe, J., Heinke, J., Gerten, D., Haddeland, I., Arnell, N. W., Clark, D. B., & Gosling, S. N. (2014). Multimodel assessment of water scarcity under climate change. Proceedings of the National Academy of Sciences, 111(9), 3245- 3250.
1.4 - Projected Change in Water Discharge (by ~2050) (rank)	92	This risk indicator is based on multi-model simulation that applies both global climate and hydrological models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP). To estimate the change at 2°C of global warming above 1980-2010 levels, simulated annual water discharge was averaged over a 31-year period with 2°C mean warming. Results are expressed in terms of relative change (%) in probability between present day (1980-2010) conditions and 2°C scenarios by 2050.	Schewe, J., Heinke, J., Gerten, D., Haddeland, I., Arnell, N. W., Clark, D. B., & Gosling, S. N. (2014). Multimodel assessment of water scarcity under climate change. Proceedings of the National Academy of Sciences, 111(9), 3245- 3250.



Indicator	Value	Description	Source
1.5 - Drought Frequency Probability (score)	2.03	This risk indicator is based on the Standardized Precipitation and Evaporation Index (SPEI). Vicente-Serrano et al. (2010) developed this multi-scalar drought index applying both precipitation and temperature data to detect, monitor and analyze different drought types and impacts in the context of global warming. The mathematical calculations used for SPEI are similar to the Standard Precipitation Index (SPI), but it has the advantage to include the role of evapotranspiration.	Vicente-Serrano, S. M., Beguería, S., & López- Moreno, J. I. (2010). A multiscalar drought index sensitive to global warming: the standardized precipitation evapotranspiration index. Journal of climate, 23(7), 1696-1718.
1.5 - Drought Frequency Probability (rank)	125	This risk indicator is based on the Standardized Precipitation and Evaporation Index (SPEI). Vicente-Serrano et al. (2010) developed this multi-scalar drought index applying both precipitation and temperature data to detect, monitor and analyze different drought types and impacts in the context of global warming. The mathematical calculations used for SPEI are similar to the Standard Precipitation Index (SPI), but it has the advantage to include the role of evapotranspiration.	Vicente-Serrano, S. M., Beguería, S., & López- Moreno, J. I. (2010). A multiscalar drought index sensitive to global warming: the standardized precipitation evapotranspiration index. Journal of climate, 23(7), 1696-1718.
1.6 - Projected Change in Drought Occurrence (by ~2050) (score)	3.00	This risk indicator is based on multi-model simulation that applies both global climate and drought models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP) . A drought threshold for pre-industrial conditions was calculated based on time-series averages. Results are expressed in terms of relative change (%) in probability between pre- industrial and 2°C scenarios.	Frieler, K., Lange, S., Piontek, F., Reyer, C. P., Schewe, J., Warszawski, L., & Geiger, T. (2017). Assessing the impacts of 1.5 C global warming–simulation protocol of the Inter- Sectoral Impact Model Intercomparison Project (ISIMIP2b). Geoscientific Model Development.
1.6 - Projected Change in Drought Occurrence (by ~2050) (rank)	134	This risk indicator is based on multi-model simulation that applies both global climate and drought models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP) . A drought threshold for pre-industrial conditions was calculated based on time-series averages. Results are expressed in terms of relative change (%) in probability between pre- industrial and 2°C scenarios.	Frieler, K., Lange, S., Piontek, F., Reyer, C. P., Schewe, J., Warszawski, L., & Geiger, T. (2017). Assessing the impacts of 1.5 C global warming–simulation protocol of the Inter- Sectoral Impact Model Intercomparison Project (ISIMIP2b). Geoscientific Model Development.
2.1 - Estimated Flood Occurrence (score)	3.03	This risk indicator is based on the recurrence of floods within the 34-year time frame period of 1985 to 2019. The occurrence of floods within a given location was estimated using data from Flood Observatory, University of Colorado. The Flood Observatory use data derived from a wide variety of news, governmental, instrumental, and remote sensing source.	Brakenridge, G. R. (2019). Global active archive of large flood events. Dartmouth Flood Observatory, University of Colorado.
2.1 - Estimated Flood Occurrence (rank)	106	This risk indicator is based on the recurrence of floods within the 34-year time frame period of 1985 to 2019. The occurrence of floods within a given location was estimated using data from Flood Observatory, University of Colorado. The Flood Observatory use data derived from a wide variety of news, governmental, instrumental, and remote sensing source.	Brakenridge, G. R. (2019). Global active archive of large flood events. Dartmouth Flood Observatory, University of Colorado.



Indicator	Value	Description	Source
2.2 - Projected Change in Flood Occurrence (by ~2050) (score)	3.00	This risk indicator is based on multi-model simulation that applies both global climate and drought models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP). The magnitude of the flood event was defined based on 100-year return period for pre-industrial conditions. Results are expressed in terms of change (%) in probability between pre- industrial and 2°C scenarios.	Frieler, K., Lange, S., Piontek, F., Reyer, C. P., Schewe, J., Warszawski, L., & Geiger, T. (2017). Assessing the impacts of 1.5 C global warming–simulation protocol of the Inter- Sectoral Impact Model Intercomparison Project (ISIMIP2b). Geoscientific Model Development.
2.2 - Projected Change in Flood Occurrence (by ~2050) (rank)	43	This risk indicator is based on multi-model simulation that applies both global climate and drought models from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP). The magnitude of the flood event was defined based on 100-year return period for pre-industrial conditions. Results are expressed in terms of change (%) in probability between pre- industrial and 2°C scenarios.	Frieler, K., Lange, S., Piontek, F., Reyer, C. P., Schewe, J., Warszawski, L., & Geiger, T. (2017). Assessing the impacts of 1.5 C global warming-simulation protocol of the Inter- Sectoral Impact Model Intercomparison Project (ISIMIP2b). Geoscientific Model Development.
3.1 - Surface Water Contamination Index (score)	4.00	The underlying data for this risk indicator is based on a broad suite of pollutants with well-documented direct or indirect negative effects on water security for both humans and freshwater biodiversity, compiled by Vörösmarty et al. (2010). The negative effects are specific to individual pollutants, ranging from impacts mediated by eutrophication such as algal blooms and oxygen depletion (e.g., caused by phosphorus and organic loading) to direct toxic effects (e.g., caused by pesticides, mercury). The overall Surface Water Contamination Index is calculated based on a range of key pollutants with different weightings according to the level of their negative effects on water security for both humans and freshwater biodiversity: soil salinization (8%), nitrogen (12%) and phosphorus (P, 13%) loading, mercury deposition (5%), pesticide loading (10%), sediment loading (17%), organic loading (as Biological Oxygen Demand, BOD; 15%), potential acidification (9%), and thermal alteration (11%).	Vörösmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P., & Davies, P. M. (2010). Global threats to human water security and river biodiversity. Nature, 467(7315), 555.



Indicator	Value	Description	Source
3.1 - Surface Water Contamination Index (rank)	28	The underlying data for this risk indicator is based on a broad suite of pollutants with well-documented direct or indirect negative effects on water security for both humans and freshwater biodiversity, compiled by Vörösmarty et al. (2010). The negative effects are specific to individual pollutants, ranging from impacts mediated by eutrophication such as algal blooms and oxygen depletion (e.g., caused by phosphorus and organic loading) to direct toxic effects (e.g., caused by pesticides, mercury). The overall Surface Water Contamination Index is calculated based on a range of key pollutants with different weightings according to the level of their negative effects on water security for both humans and freshwater biodiversity: soil salinization (8%), nitrogen (12%) and phosphorus (P, 13%) loading, mercury deposition (5%), pesticide loading (10%), sediment loading (17%), organic loading (as Biological Oxygen Demand, BOD; 15%),	Vörösmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P., & Davies, P. M. (2010). Global threats to human water security and river biodiversity. Nature, 467(7315), 555.
4.1 - Fragmentation Status of Rivers (score)	4.03	potential acidification (9%), and thermal alteration (11%). This risk indicator is based on the data set by Grill et al. (2019) mapping the world's free-flowing rivers. Grill et al. (2019) compiled a geometric network of the global river system and associated attributes, such as hydro-geometric properties, as well as pressure indicators to calculate an integrated connectivity status index (CSI). While only rivers with high levels of connectivity in their entire length are classified as free-flowing, rivers of CSI < 95% are considered as fragmented at a certain degree.	Grill, G., Lehner, B., Thieme, M., Geenen, B., Tickner, D., Antonelli, F., & Macedo, H. E. (2019). Mapping the world's free-flowing rivers. Nature, 569(7755), 215.
4.1 - Fragmentation Status of Rivers (rank)	13	This risk indicator is based on the data set by Grill et al. (2019) mapping the world's free-flowing rivers. Grill et al. (2019) compiled a geometric network of the global river system and associated attributes, such as hydro-geometric properties, as well as pressure indicators to calculate an integrated connectivity status index (CSI). While only rivers with high levels of connectivity in their entire length are classified as free-flowing, rivers of CSI < 95% are considered as fragmented at a certain degree.	Grill, G., Lehner, B., Thieme, M., Geenen, B., Tickner, D., Antonelli, F., & Macedo, H. E. (2019). Mapping the world's free-flowing rivers. Nature, 569(7755), 215.
4.2 - Catchment Ecosystem Services Degradation Level (tree cover loss) (score)	3.46	For this risk indicator, tree cover loss was applied as a proxy to represent catchment ecosystem services degradation since forests play an important role in terms of water regulation, supply and pollution control. The forest cover data is based on Hansen et al.'s global Landsat data at a 30-meter spatial resolution to characterize forest cover and change. The authors defined trees as vegetation taller than 5 meters in height, and forest cover loss as a stand-replacement disturbance, or a change from a forest to non-forest state, during the period 2000 – 2018.	Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A. A., Tyukavina, A., & Kommareddy, A. (2013). High-resolution global maps of 21st-century forest cover change. science, 342(6160), 850-853.



Indicator	Value	Description	Source
4.2 - Catchment Ecosystem Services Degradation Level (tree cover loss) (rank)	32	For this risk indicator, tree cover loss was applied as a proxy to represent catchment ecosystem services degradation since forests play an important role in terms of water regulation, supply and pollution control. The forest cover data is based on Hansen et al.'s global Landsat data at a 30-meter spatial resolution to characterize forest cover and change. The authors defined trees as vegetation taller than 5 meters in height, and forest cover loss as a stand-replacement disturbance, or a change from a forest to non-forest state, during the period 2000 – 2018.	Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A. A., Tyukavina, A., & Kommareddy, A. (2013). High-resolution global maps of 21st-century forest cover change. science, 342(6160), 850-853.
4.3 - Projected Impacts on Freshwater Biodiversity (score)	2.00	The study by Tedesco et al. (2013) to project changes [% increase or decrease] in extinction rate by ~2090 of freshwater fish due to water availability loss from climate change is used as a proxy to estimate the projected impacts on freshwater biodiversity.	Tedesco, P. A., Oberdorff, T., Cornu, J. F., Beauchard, O., Brosse, S., Dürr, H. H., & Hugueny, B. (2013). A scenario for impacts of water availability loss due to climate change on riverine fish extinction rates. Journal of Applied Ecology, 50(5), 1105-1115.
4.3 - Projected Impacts on Freshwater Biodiversity (rank)	144	The study by Tedesco et al. (2013) to project changes [% increase or decrease] in extinction rate by ~2090 of freshwater fish due to water availability loss from climate change is used as a proxy to estimate the projected impacts on freshwater biodiversity.	Tedesco, P. A., Oberdorff, T., Cornu, J. F., Beauchard, O., Brosse, S., Dürr, H. H., & Hugueny, B. (2013). A scenario for impacts of water availability loss due to climate change on riverine fish extinction rates. Journal of Applied Ecology, 50(5), 1105-1115.
5.1 - Freshwater Policy Status (SDG 6.5.1) (score)	1.00	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "National Water Resources Policy" indicator, which corresponds to one of the three national level indicators under the Enabling Environment category.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
5.1 - Freshwater Policy Status (SDG 6.5.1) (rank)	183	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "National Water Resources Policy" indicator, which corresponds to one of the three national level indicators under the Enabling Environment category.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
5.2 - Freshwater Law Status (SDG 6.5.1) (score)	1.00	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "National Water Resources Law(s)" indicator, which corresponds to one of the three national level indicators under the Enabling Environment category. For SDG 6.5.1, enabling environment depicts the conditions that help to support the implementation of IWRM, which includes legal and strategic planning tools for IWRM.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.



Indicator	Value	Description	Source
5.2 - Freshwater Law Status (SDG 6.5.1) (rank)	176	 This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "National Water Resources Law(s)" indicator, which corresponds to one of the three national level indicators under the Enabling Environment category. For SDG 6.5.1, enabling environment depicts the conditions that help to support the implementation of IWRM, which includes legal and strategic planning tools for IWRM. 	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
5.3 - Implementation Status of Water Management Plans (SDG 6.5.1) (score)	2.00	 This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "National IWRM plans" indicator, which corresponds to one of the three national level indicators under the Enabling Environment category. For SDG 6.5.1, enabling environment depicts the conditions that help to support the implementation of IWRM, which includes legal and strategic planning tools for IWRM. 	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
5.3 - Implementation Status of Water Management Plans (SDG 6.5.1) (rank)	160	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "National IWRM plans" indicator, which corresponds to one of the three national level indicators under the Enabling Environment category. For SDG 6.5.1, enabling environment depicts the conditions that help to support the implementation of IWRM, which includes legal and strategic planning tools for IWRM.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
6.1 - Corruption Perceptions Index (score)	1.00	This risk Indicator is based on the latest Transparency International's data: the Corruption Perceptions Index 2018. This index aggregates data from a number of different sources that provide perceptions of business people and country experts on the level of corruption in the public sector.	Transparency International (2019). Corruption Perceptions Index 2018. Berlin: Transparency International.
6.1 - Corruption Perceptions Index (rank)	185	This risk Indicator is based on the latest Transparency International's data: the Corruption Perceptions Index 2018. This index aggregates data from a number of different sources that provide perceptions of business people and country experts on the level of corruption in the public sector.	Transparency International (2019). Corruption Perceptions Index 2018. Berlin: Transparency International.
6.2 - Freedom in the World Index (score)	1.00	This risk indicator is based on Freedom House (2019), an annual global report on political rights and civil liberties, composed of numerical ratings and descriptive texts for each country and a select group of territories. The 2019 edition involved more than 100 analysts and more than 30 advisers with global, regional, and issue-based expertise to covers developments in 195 countries and 14 territories from January 1, 2018, through December 31, 2018.	Freedom House (2019). Freedom in the world 2019. Washington, DC: Freedom House.



Indicator	Value	Description	Source
6.2 - Freedom in the World Index(rank)	170	This risk indicator is based on Freedom House (2019), an annual global report on political rights and civil liberties, composed of numerical ratings and descriptive texts for each country and a select group of territories. The 2019 edition involved more than 100 analysts and more than 30 advisers with global, regional, and issue-based expertise to covers developments in 195 countries and 14 territories from January 1, 2018, through December 31, 2018.	Freedom House (2019). Freedom in the world 2019. Washington, DC: Freedom House.
6.3 - Business Participation in Water Management (SDG 6.5.1) (score)	1.00	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "Business Participation in Water Resources Development, Management and Use" indicator, which corresponds to one of the six national level indicators under the Institutions and Participation category.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
6.3 - Business Participation in Water Management (SDG 6.5.1) (rank)	178	This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "Business Participation in Water Resources Development, Management and Use" indicator, which corresponds to one of the six national level indicators under the Institutions and Participation category.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
7.1 - Management Instruments for Water Management (SDG 6.5.1) (score)	1.00	 This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "Sustainable and efficient water use management" indicator, which corresponds to one of the five national level indicators under the Management Instruments category. For SDG 6.5.1, management instruments refer to the tools and activities that enable decision-makers and users to make rational and informed choices between alternative actions. 	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
7.1 - Management Instruments for Water Management (SDG 6.5.1) (rank)	178	 This risk indicator is based on SDG 6.5.1. Degree of IWRM Implementation "Sustainable and efficient water use management" indicator, which corresponds to one of the five national level indicators under the Management Instruments category. For SDG 6.5.1, management instruments refer to the tools and activities that enable decision-makers and users to make rational and informed choices between alternative actions. 	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.



Indicator	Value	Description	Source
7.2 - Groundwater Monitoring Data Availability and Management (score)	1.00	This risk indicator is based on the data set by UN IGRAC (2019) to determine the level of availability of groundwater monitoring data at country level as groundwater management decisions rely strongly on data availability. The level of groundwater monitoring data availability for groundwater management is determined according to a combination of three criteria developed by WWF and IGRAC: 1) Status of country groundwater monitoring programme, 2) groundwater data availability for NGOs and 3) Public access to processed groundwater monitoring data.	UN IGRAC (2019). Global Groundwater Monitoring Network GGMN Portal. UN International Groundwater Resources Assessment Centre (IGRAC).
7.2 - Groundwater Monitoring Data Availability and Management (rank)	177	This risk indicator is based on the data set by UN IGRAC (2019) to determine the level of availability of groundwater monitoring data at country level as groundwater management decisions rely strongly on data availability. The level of groundwater monitoring data availability for groundwater management is determined according to a combination of three criteria developed by WWF and IGRAC: 1) Status of country groundwater monitoring programme, 2) groundwater data availability for NGOs and 3) Public access to processed groundwater monitoring data.	UN IGRAC (2019). Global Groundwater Monitoring Network GGMN Portal. UN International Groundwater Resources Assessment Centre (IGRAC).
7.3 - Density of Runoff Monitoring Stations (score)	2.00	The density of monitoring stations for water quantity was applied as proxy to develop this risk indicator. The Global Runoff Data Base was used to estimate the number of monitoring stations per 1000km2 of the main river system (data base access date: May 2018).	BfG (2019). Global Runoff Data Base. German Federal Institute of Hydrology (BfG).
7.3 - Density of Runoff Monitoring Stations (rank)	184	The density of monitoring stations for water quantity was applied as proxy to develop this risk indicator. The Global Runoff Data Base was used to estimate the number of monitoring stations per 1000km2 of the main river system (data base access date: May 2018).	BfG (2019). Global Runoff Data Base. German Federal Institute of Hydrology (BfG).
8.1 - Access to Safe Drinking Water (score)	1.00	This risk indicator is based on the Joint Monitoring Programme for Water Supply, Sanitation and Hygiene (UNICEF/WHO) 2019 data. It provides estimates on the use of water, sanitation and hygiene by country for the period 2000-2017.	WHO & UNICEF (2019). Estimates on the use of water, sanitation and hygiene by country (2000- 2017). Joint Monitoring Programme for Water Supply, Sanitation and Hygiene.
8.1 - Access to Safe Drinking Water (rank)	165	This risk indicator is based on the Joint Monitoring Programme for Water Supply, Sanitation and Hygiene (UNICEF/WHO) 2019 data. It provides estimates on the use of water, sanitation and hygiene by country for the period 2000-2017.	WHO & UNICEF (2019). Estimates on the use of water, sanitation and hygiene by country (2000- 2017). Joint Monitoring Programme for Water Supply, Sanitation and Hygiene.
8.2 - Access to Sanitation (score)	1.00	This risk indicator is based on the Joint Monitoring Programme for Water Supply, Sanitation and Hygiene (UNICEF/WHO) 2019 data. It provides estimates on the use of water, sanitation and hygiene by country for the period 2000-2017.	WHO & UNICEF (2019). Estimates on the use of water, sanitation and hygiene by country (2000- 2017). Joint Monitoring Programme for Water Supply, Sanitation and Hygiene.



Indicator	Value	Description	Source
8.2 - Access to Sanitation (rank)	174	This risk indicator is based on the Joint Monitoring Programme for Water Supply, Sanitation and Hygiene (UNICEF/WHO) 2019 data. It provides estimates on the use of water, sanitation and hygiene by country for the period 2000-2017.	WHO & UNICEF (2019). Estimates on the use of water, sanitation and hygiene by country (2000- 2017). Joint Monitoring Programme for Water Supply, Sanitation and Hygiene.
8.3 - Financing for Water Resource Development and Management (SDG 6.5.1) (score)	1.00	This risk indicator is based on the average 'Financing' score of UN SDG 6.5.1. Degree of IWRM Implementation database. UN SDG 6.5.1 database contains a category on financing which assesses different aspects related to budgeting and financing made available and used for water resources development and management from various sources.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
8.3 - Financing for Water Resource Development and Management (SDG 6.5.1) (rank)	184	This risk indicator is based on the average 'Financing' score of UN SDG 6.5.1. Degree of IWRM Implementation database. UN SDG 6.5.1 database contains a category on financing which assesses different aspects related to budgeting and financing made available and used for water resources development and management from various sources.	UN Environment (2018). Progress on integrated water resources management. Global baseline for SDG 6 Indicator 6.5.1: degree of IWRM implementation.
9.1 - Cultural Diversity (score)	3.00	Water is a social and cultural good. The cultural diversity risk indicator was included in order to acknowledge that businesses face reputational risk due to the importance of freshwater for indigenous and traditional people in their daily life, religion and culture. This risk indicator is based on Oviedo and Larsen (2000) data set, which mapped the world's ethnolinguistic groups onto the WWF map of the world's ecoregions. This cross-mapping showed for the very first time the significant overlap that exists between the global geographic distribution of biodiversity and that of linguistic diversity.	Oviedo, G., Maffi, L., & Larsen, P. B. (2000). Indigenous and traditional peoples of the world and ecoregion conservation: An integrated approach to conserving the world's biological and cultural diversity. Gland: WWF (World Wide Fund for Nature) International.
9.1 - Cultural Diversity (rank)	75	Water is a social and cultural good. The cultural diversity risk indicator was included in order to acknowledge that businesses face reputational risk due to the importance of freshwater for indigenous and traditional people in their daily life, religion and culture. This risk indicator is based on Oviedo and Larsen (2000) data set, which mapped the world's ethnolinguistic groups onto the WWF map of the world's ecoregions. This cross-mapping showed for the very first time the significant overlap that exists between the global geographic distribution of biodiversity and that of linguistic diversity.	Oviedo, G., Maffi, L., & Larsen, P. B. (2000). Indigenous and traditional peoples of the world and ecoregion conservation: An integrated approach to conserving the world's biological and cultural diversity. Gland: WWF (World Wide Fund for Nature) International.
10.1 - Freshwater Endemism (score)	5.00	The underlying data set for this risk indicator comes from the Freshwater Ecoregions of the World (FEOW) 2015 data developed by WWF and TNC. Companies operating in basins with higher number of endemic fish species are exposed to higher reputational risks.	WWF & TNC (2015). Freshwater Ecoregions of the World.



Indicator	Value	Description	Source
10.1 - Freshwater Endemism (rank)	16	The underlying data set for this risk indicator comes from the Freshwater Ecoregions of the World (FEOW) 2015 data developed by WWF and TNC. Companies operating in basins with higher number of endemic fish species are exposed to higher reputational risks.	WWF & TNC (2015). Freshwater Ecoregions of the World.
10.2 - Freshwater Biodiversity Richness (score)	4.00	The underlying data set for this risk indicator comes from the Freshwater Ecoregions of the World (FEOW) 2015 data developed by WWF and TNC. Count of fish species is used as a representation of freshwater biodiversity richness. Companies operating in basins with higher number of fish species are exposed to higher reputational risks.	WWF & TNC (2015). Freshwater Ecoregions of the World.
10.2 - Freshwater Biodiversity Richness (rank)	60	The underlying data set for this risk indicator comes from the Freshwater Ecoregions of the World (FEOW) 2015 data developed by WWF and TNC. Count of fish species is used as a representation of freshwater biodiversity richness. Companies operating in basins with higher number of fish species are exposed to higher reputational risks.	WWF & TNC (2015). Freshwater Ecoregions of the World.
11.1 - National Media Coverage (score)	3.00	This risk indicator is based on joint qualitative research by WWF and Tecnoma (Typsa Group). It indicates how aware local residents typically are of water-related issues due to national media coverage. The status of the river basin (e.g., scarcity and pollution) is taken into account, as well as the importance of water for livelihoods (e.g., food and shelter).	WWF & Tecnoma (TYPSA Group)
11.1 - National Media Coverage (rank)	160	This risk indicator is based on joint qualitative research by WWF and Tecnoma (Typsa Group). It indicates how aware local residents typically are of water-related issues due to national media coverage. The status of the river basin (e.g., scarcity and pollution) is taken into account, as well as the importance of water for livelihoods (e.g., food and shelter).	WWF & Tecnoma (TYPSA Group)
11.2 - Global Media Coverage (score)	2.00	This risk indicator is based on joint qualitative research by WWF and Tecnoma (Typsa Group). It indicates how aware people are of water- related issues due to global media coverage. Familiarity to and media coverage of the region and regional water-related disasters are taken into account.	WWF & Tecnoma (TYPSA Group)
11.2 - Global Media Coverage (rank)	137	This risk indicator is based on joint qualitative research by WWF and Tecnoma (Typsa Group). It indicates how aware people are of water- related issues due to global media coverage. Familiarity to and media coverage of the region and regional water-related disasters are taken into account.	WWF & Tecnoma (TYPSA Group)



Indicator	Value	Description	Source
12.1 - Conflict News Events (RepRisk) (score)	3.00	This risk indicator is based on 2018 data collected by RepRisk on counts and registers of documented negative incidents, criticism and controversies that can affect a company's reputational risk. These negative news events are labelled per country and industry class.	RepRisk & WWF (2019). Due diligence database on ESG and business conduct risks. RepRisk.
12.1 - Conflict News Events (RepRisk) (rank)	100	This risk indicator is based on 2018 data collected by RepRisk on counts and registers of documented negative incidents, criticism and controversies that can affect a company's reputational risk. These negative news events are labelled per country and industry class.	RepRisk & WWF (2019). Due diligence database on ESG and business conduct risks. RepRisk.
12.2 - Hydro-political Risk (score)	1.18	This risk indicator is based on the assessment of hydro-political risk by Farinosi et al. (2018). More specifically, it is based on the results of spatial modelling by Farinosi et al. (2018) that determined the main parameters affecting water cross-border conflicts and calculated the likelihood of hydro-political issues.	Farinosi, F., Giupponi, C., Reynaud, A., Ceccherini, G., Carmona-Moreno, C., De Roo, A., & Bidoglio, G. (2018). An innovative approach to the assessment of hydro-political risk: A spatially explicit, data driven indicator of hydro- political issues. Global environmental change, 52, 286-313.
12.2 - Hydro-political Risk (rank)	183	This risk indicator is based on the assessment of hydro-political risk by Farinosi et al. (2018). More specifically, it is based on the results of spatial modelling by Farinosi et al. (2018) that determined the main parameters affecting water cross-border conflicts and calculated the likelihood of hydro-political issues.	Farinosi, F., Giupponi, C., Reynaud, A., Ceccherini, G., Carmona-Moreno, C., De Roo, A., & Bidoglio, G. (2018). An innovative approach to the assessment of hydro-political risk: A spatially explicit, data driven indicator of hydro- political issues. Global environmental change, 52, 286-313.
Population, total (#)	582972	Population, total	The World Bank 2018, Data , hompage accessed 20/04/2018
GDP (current US\$)	58631324559	GDP (current US\$)	The World Bank 2018, Data , hompage accessed 20/04/2018
EPI 2018 score (0-100)	79.12	Environmental Performance Index	
WGI -Voice and Accountability (0-100)	97.62	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, The Worldwide Governance Indicators: Methodology and Analytical Issues (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132



Indicator	Value	Description	Source
WGI -Political stability no violence (0-100)	96.55	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, The Worldwide Governance Indicators: Methodology and Analytical Issues (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132
WGl - Government Effectiveness (0-100)	93.27	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, The Worldwide Governance Indicators: Methodology and Analytical Issues (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132
WGI - Regulatory Quality (0-100)	93.75	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, The Worldwide Governance Indicators: Methodology and Analytical Issues (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132
WGI - Rule of Law (0-100)	93.75	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, The Worldwide Governance Indicators: Methodology and Analytical Issues (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132
WGI - Control of Corruption (0-100)	97.60	Water Governance Indicator	Kaufmann, Daniel and Kraay, Aart and Mastruzzi, Massimo, The Worldwide Governance Indicators: Methodology and Analytical Issues (September 2010). World Bank Policy Research Working Paper No. 5430. Available at SSRN: https://ssrn.com/abstract=1682132



Indicator	Value	Description	Source
WRI BWS all industries (0-5)	2.51	WRI Baseline Water Stress (BWS)	Gassert, F., P. Reig, T. Luo, and A. Maddocks. 2013. "Aqueduct country and river basin rankings: a weighted aggregation of spatially distinct hydrological indicators." Working paper. Washington, DC: World Resources Institute, December 2013. Available online at http://wri.org/publication/aqueduct-country- river-basin-rankings.
WRI BWS Ranking (1=very high)	77	WRI Baseline Water Stress (BWS)	Gassert, F., P. Reig, T. Luo, and A. Maddocks. 2013. "Aqueduct country and river basin rankings: a weighted aggregation of spatially distinct hydrological indicators." Working paper. Washington, DC: World Resources Institute, December 2013. Available online at http://wri.org/publication/aqueduct-country- river-basin-rankings.
Baseline Water Stress (BWS) - 2020 BAU (1=very high)	55	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct- projected-water-stress-country-rankings.
Baseline Water Stress (BWS) - 2020 Optimistic (increasing rank describes lower risk)	54	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct- projected-water-stress-country-rankings.
Baseline Water Stress (BWS) - 2020 Pessimistic (increasing rank describes lower risk)	54	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct- projected-water-stress-country-rankings.



Indicator	Value	Description	Source
Baseline Water Stress (BWS) - 2030 BAU (increasing rank describes lower risk)	59	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct- projected-water-stress-country-rankings.
Baseline Water Stress (BWS) - 2030 Optimistic (increasing rank describes lower risk)	58	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct- projected-water-stress-country-rankings.
Baseline Water Stress (BWS) - 2030 Pessimistic (increasing rank describes lower risk)	59	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct- projected-water-stress-country-rankings.
Baseline Water Stress (BWS) - 2040 BAU (increasing rank describes lower risk)	63	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct- projected-water-stress-country-rankings.
Baseline Water Stress (BWS) - 2040 Optimistic (increasing rank describes lower risk)	60	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct- projected-water-stress-country-rankings.
Baseline Water Stress (BWS) - 2040 Pessimistic (increasing rank describes lower risk)	64	WRI country ranking	Luo, T., R. Young, and P. Reig. 2015. "Aqueduct projected water stress rankings." Technical note. Washington, DC: World Resources Institute, August 215. Available online at http://www.wri.org/publication/aqueduct- projected-water-stress-country-rankings.



Indicator	Value	Description	Source
Total water footprint of national consumption (m3/a/cap)	2514.30	WFN Water Footprint Data	Mekonnen, M.M. and Hoekstra, A.Y. (2011) National water footprint accounts: The green, blue and grey water footprint of production and consumption, Value of Water Research Report Series No. 50, UNESCO-IHE, Delft, the Netherlands.http://www.waterfootprint.org/Rep orts/Report50-NationalWaterFootprints-Vol1.pdf
Ratio external / total water footprint (%)	75.09	WFN Water Footprint Data	Mekonnen, M.M. and Hoekstra, A.Y. (2011) National water footprint accounts: The green, blue and grey water footprint of production and consumption, Value of Water Research Report Series No. 50, UNESCO-IHE, Delft, the Netherlands.http://www.waterfootprint.org/Rep orts/Report50-NationalWaterFootprints-Vol1.pdf
Area equipped for full control irrigation: total (1000 ha)	0.00	Aquastat - Irrigation	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Area equipped for irrigation: total (1000 ha)	0.00	Aquastat - Irrigation	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
% of the area equipped for irrigation actually irrigated (%)	100.00	Aquastat - Irrigation	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Electricity production from hydroelectric sources (% of total)	7.51	World Development Indicators	The World Bank 2018, Data , hompage accessed 20/04/2018
Total internal renewable water resources (IRWR) (10^9 m3/year)	1.00	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Total internal renewable water resources (IRWR) (10^9 m3/year)	2.50	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Water resources: total external renewable (10^9 m3/year)	1.00	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13



Indicator	Value	Description	Source
Total renewable water resources (10^9 m3/year)	3.50	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Dependency ratio (%)	71.43	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
Total renewable water resources per capita (m3/inhab/year)	6172.00	Aquastat - Water Ressources	FAO. 2016. AQUASTAT website. Food and Agriculture Organization of the United Nations (FAO). Website accessed on 2018/04/13
World happiness [0-8]	6.91	WorldHappinessReport.org	World Happiness Report, homepage accessed 20/04/2018



Country Aspects

1. PHYSICAL ASPECTS

1.1.WATER RESOURCES

1.1.1.WATER RESOURCES

Luxembourg has a moderate oceanic western European climate with mild winters and comfortable summers. Absolute minimum and maximum air temperatures in the reference period 1961-1990 range from –17.8°C in January (1979) to 35.1°C in July (1964).

The long-term average of annual precipitation is 934 mm/year. The long-term average of annual renewable water resources is stated as 3,100 million m³/year of which 32 per cent are considered to be internal water resources (1,000 million m³) and as much as 68 per cent are considered to be external water resources (2,100 million of m³). As at 2009 the total dam (reservoir) capacity was stated to be 62 million m³.

Luxembourg has two river basin districts (RBDs) which are the Maas and the Rhine. Both are international, sharing water courses with France in the southeast, Belgium in the northeast and Germany in the west. Consultation on draft river basin management plans took place between 22 December 2008 and 30 June 2009. River basin management plans for Luxembourg were adopted in December 2009.

1.1.2.WATER USE

The usage intensity of available resources in households, industry and agriculture is relatively low compared to other developed countries, reflecting the low level of abstraction for agriculture. Daily household consumption is 140 litres per capita. There is little loss through leakage and many of the water supply systems have been upgraded over the last 10 years.

Water withdrawal in the country has been estimated at a total of 69.2 million m³. This water abstraction can be itemized by user sector. It was estimated that 47 million m³ was dedicated to urban use in 2009 (around 67 per cent of total water consumption). In 1999, water withdrawal for agriculture was estimated at only 0.2 million m³; and for industrial use, 22 million m³ (which represents around 32 per cent of the total abstraction).

Groundwater provides 57 per cent of the 47 million m3 of drinking water used annually in Luxembourg. Also, in the southern industrial region, industries and utilities must draw from the water table of the Luxembourg sandstone aquifer, located in the centre of the country, and from the Esch-sur-Sûre reservoir in the north to cover their water needs.

Under the legislation of Luxembourg regarding water (The Water Act), there must be a balance between depletion and renewal of underground water sources so that they will be in good condition by no later than 2015. The preference given to surface tapping over groundwater

pumping eliminates the risk of over-exploitation of the sandstone aquifer, which supplies more than half of the country's drinking water. On the other hand, the Esch-sur-Sûre reservoir, which provides 43 per cent of the water supply to public utilities in Luxembourg, is in a critical state of eutrophication.

While the demand for water from industry has decreased with the improvement of industrial processes, notably in the metallurgy sector, household consumption has increased by 1.35 per cent per year over the last 15 years, reflecting the country's strong demographic growth and the steady increase in cross-border workers.

1.2. WATER QUALITY, ECOSYSTEMS AND HUMAN HEALTH

It is estimated that only 30 per cent of surface water bodies will comply with the EU's 2015 targets for chemical and biological quality as determined under the EU Water Framework Directive (WFD) (Directive 2000/60/EC). The continuous significant demographic growth that Luxembourg has experienced these last 25 years - and that will most likely continue in future - is the main threat to water resources and water quality. With regard to drinking water, sources have not yet been protected, despite a legal obligation to do so dating back more than 15 years. Moreover, rural development policies have focused more on farm modernization and the continued use of agricultural land than on the targeted protection of water resources.

According to the Water Act, all surface water bodies must be protected, improved or restored to meet the definition of "good status" by the end of 2015. However, it is estimated that at least 72 per cent of surface water bodies (watercourses and reservoirs) will not meet the 2015 chemical and biological quality targets under the WFD. While the pollution level in watercourses has decreased slightly in recent years, 39 per cent of watercourses are still heavily polluted and 54 per cent moderately polluted.

With regard to new contaminants, the watercourses of the Alzette and the Mess, located in industrialised and heavily populated environments, have been found to contain xenobiotic pollution from antibiotics, analgesics and hormones. There are several xenobiotic pollutants coming from diffuse pollution that are found in all major watercourses. These pollutants can not be eliminated by the existing purification plants.

The quality of all drinking water resources (250 catchment sources, 50 drilling holes and the Eschsur-Sûre reservoir) is regularly monitored. Suppliers are responsible for monitoring the quality of the water they deliver for human consumption. The Grand Ducal Regulation of 7 October 2002, which implements Directive 98/83/EC on the quality of water intended for human consumption, requires drinking water suppliers to audit their infrastructure and assess the state of water resources. A total of 97 of the 116 municipalities and the seven inter-municipal syndicates had finished their audit by the end of March 2010.

Under the terms of the Water Act, all bodies of groundwater must be protected, improved and



restored to "good status" by the end of 2015. The main groundwater pollutants are nitrates and pesticides. At a national level, the nitrogen content has decreased considerably since the early 1990s, dropping from 200kg N/hectare to 111 kg N/hectare in 2004 (last available year). However, a recent study showed that 40 per cent of the surface area that drains into drinking water sources discharges water containing 25-50mg/litre of nitrates. Moreover, some of the sources show a clear trend towards deterioration. At a national level, half of the nitrogen input comes from the use of chemical fertilizers, and a third from livestock effluents; the rest is atmospherically deposited. In up to 90 per cent of the monitored groundwater sampling sites pesticides are detected, sometimes at concentrations that exceed the threshold value of 100mg/litre. This indicates that not only is the Luxembourg sandstone aquifer more vulnerable to pollution than the aquifers of neighbouring regions, but also that there is a lack of protection for the abstraction areas.

2. GOVERNANCE ASPECTS

All the environmental challenges are considered by the authorities through a set of plans including a cross-sector integrated plan (IVL) (Integratives Verkehrs und Landesentwicklungskonzept für Luxemburg), various sectoral Action Plans – some of them deriving from the cross-sector integrated plan – and Action Plans aiming at fulfilling the objective set for Luxembourg by the EU 'Climate and Energy Package'.

2.1.WATER INSTITUTIONS

In 1999, the Government of Luxembourg opted for a comprehensive water management policy, designed to consolidate the various aspects of the water economy and create a tool for integrated water management. The Water Management Administration, which reports to the Minister for the Interior, was created in 2004, amalgamating the various units responsible for water protection and management that had previously operated within other administrations.

2.2.WATER MANAGEMENT

The Water Act has been implemented to consolidate water legislation and give effect to the WFD and the EU Floods Directive (2007/60/EC) in national law. It also called for management plans to be prepared for Luxembourg's two river basins and these were made public at the end of 2009. Furthermore, it requires the establishment of a "general municipal plan for the urban water cycle" in each municipality. This plan must contain an inventory of underground waters, water supply and sanitation infrastructure; areas listed in the protected zones registry, and details of flood-prone districts. These municipal plans will provide the basis for preparing a national urban water cycle plan.

One particular element that is being implemented in Luxembourg is the structure of water pricing and the introduction of the principle of full cost recovery for drinking water supply and urban sewage treatment. In addition to water supply and sanitation service charges, which are levied by the service providers, the law introduces an abstraction tax and a pollution tax, income from which goes to the Water Management Fund. The abstraction tax is levied on anyone who draws surface water or groundwater, and is based on the volume of water drawn (measured by a metering device installed by the user). In addition to the public utilities, which provide 43 million m3 of water annually (70 per cent for the public network and 30 per cent for industry), the agri-food industry extracts an additional 4 million m3 of groundwater, but not all these extraction sources are equipped with metering devices.

The discharge of waste water into surface or underground water sources is subject to a pollution tax. The tax is proportionate to the units of pollutant load in the water discharged. It must be paid when any of the following thresholds are exceeded: 250kg/year for chemical oxygen demand (COD); 125kg/year for nitrogen (N); 15kg/year for phosphorus (P); or 5.2kg/year for suspended particulate matter (SPM). The volume of water discharged is equal to the volume of water drawn from the public distribution network. A 10-20 per cent reduction in the tax is offered to municipalities that have installed rainwater treatment and management facilities in their network. For industry, the number of units of pollutant load taken as a basis for calculating the tax is the authorised pollutant load. However, if that load is exceeded, the tax may be increased. It can also be reduced if the pollutant load is at least 20 per cent less than what would result from the discharge authorisation.

The Water Management Fund was created in 1999 to subsidise sewage treatment and is financed by budgetary allocations. The Fund is also considering resorting to loans from the European Investment Bank, so as not to hinder the development of sanitation and waste water treatment infrastructure in the coming years. The Fund can cover up to 90 per cent of the municipality's capital costs for sewerage and sewage treatment. The Water Act expands the scope of the Fund. It authorises coverage of: (i) up to 50 per cent for measures to protect water resources intended for human consumption (with the exception of agricultural activity); (ii) up to 80 per cent of the cost of flood risk abatement; and (iii) up to 100 per cent of watercourse rehabilitation costs.

The water related law also allows the Fund to cover up to 100 per cent of expenditure on projects of national interest to safeguard the quality of surface and groundwater or protect available water resources in the long term. The Water Management Fund must distribute its revenues on the basis of a cost-benefit analysis for the projects selected. The Fund itself is administered by a management committee, including representatives of the ministries responsible for water management, the budget, agriculture, health and the environment.

Financial assistance to the municipalities from the Water Management Fund has been doubled to help them cover 90 per cent of investments in sewerage and sewage treatment. However, although the proportion of the population connected to a waste water treatment plant is very high (95 per cent), only 22 per cent is connected to a tertiary treatment station, even though the entire country is classified as a sensitive area under the EU Urban Waste Water Treatment Directive (91/271/EEC). A dual-channel system to separate rainwater (which can re-infiltrate the water table naturally) and sewage (which requires purification) is still not in place, with the exception of the cities of Luxembourg and Esch-sur-Alzette and in new housing developments.

In recent decades, Luxembourg has suffered numerous bouts of flooding; in 1983 along the Moselle and in 1993, 1995 and 2003 in the Sûre basin. Since 1995, the government has been



covering 50 per cent of the cost of flood control measures. Nevertheless, according to the the Water Act, the Water Management Administration is to work with the municipalities and administrations concerned to establish a master plan for flood risk management, reflecting the objectives of the EU Floods Directive.

Between 1998 and 2002, the Spatial Planning Department established a partial management plan for flood zones and retention zones for various communities affected by high water along the Moselle and its tributaries. During the period 2002-09, the project was upgraded to a Flood Vulnerability Atlas and the investigated area has been raised to 15 of Luxembourg's watercourses. The results of the TIMIS flood project (Transnational Internet Map Information System on Flooding) were posted on the internet.

In application of article 13 of the EU Floods Directive, flood hazard maps and flood risk maps were to be completed by the end of 2010 and sent out to the municipalities for validation. The mapping of flood zones and flood risks is to be superimposed onto the municipalities' general land use plans. In particular, new urban development must be prohibited in flood zones, unless the retention volume lost can be offset and the development does not increase risks upstream and downstream.

Beyond conventional dike-building measures, the flood risk management plans will focus on improving the eco-morphological structure of riverbeds and restoring natural water retention areas. The costs are borne by the municipalities; however they may receive government subsidies of up to 50 per cent (or 80 per cent for inter-municipal works). Agricultural restrictions can also be imposed with a view to limiting the leaching of pollutants. For example, surface water management seeks to keep watercourses flowing freely and to maintain their banks in good condition. Maintenance focuses on riverbeds and the vegetation of river banks, riparian zones and floodplains. Up to 50 per cent of the costs incurred are borne by the government. Another example is the rehabilitation of watercourses and associated wetlands to enhance their flood control function. A number of "watershed level - are now underway. The cost of rehabilitation is borne by the municipalities, although they may receive state subsidies of up to 100 per cent.

Achieving sound hydro-morphological status for watercourses, as required under the Water Act, will largely depend on cooperation with riparian owners to maintain vegetation along river banks and/or state purchase of the river banks. Several "flood partnerships" (national and transnational) have been founded in order to maintain a certain level of flood risk awareness among the population and to involve the public in the implementation of the EU Floods Directive. Concerted action with neighbouring countries has led to the implementation of the Flood Action Plan Moselle since 1998 (with Germany, Belgium and France), under the aegis of the International Commissions for the Protection of the Moselle and the Sarre; and to closer cooperation on flood risk information (with France and Germany). The Flood Action Plan Moselle provides a basis for establishing the flood risk management plan in conformity with the Floods Directive.

The main Directives and related legislation at the European level, regarding the domain of water resources management are:

•Directive 80/68/EEC of 17 December, 1979 on the protection of groundwater against pollution caused by certain dangerous substances;

•Directive 82/176/EEC of 22 March, 1982 on limit values and quality objectives for mercury discharges by the chlor-alkali electrolysis industry; covers inland surface water, territorial waters and internal coastal waters;

•Directive 83/513/EEC of 26 September, 1983 on limit values and quality objectives for cadmium discharges; sets limit values and quality objectives for cadmium discharges in the aquatic environment;

•Directive 84/156/EEC of 8 March, 1984 on limit values and quality objectives for mercury discharges by sectors other than the chlor-alkali electrolysis industry; sets limit values and quality objectives for mercury discharges in sectors other than the chlor-alkali electrolysis industry;

•Directive 84/491/EEC of 9 October, 1984 on limit values and quality objectives for discharges of hexachlorocyclohexane;

•Directive 91/271/EEC of 21 May, 1991 concerning urban waste water treatment;

•Directive 91/676/EEC of 12 December, 1991 concerning the protection of waters against pollution caused by nitrates from agricultural sources (known as the Nitrates Directive);

•Directive 98/83/EC of 3 November, 1998 on the quality of water;

•Directive 2000/60/EC of the European Parliament and of the Council establishing a framework for the Community action in the field of water policy;

•Regulation (EC) No 648/2004 of the European Parliament and of the Council of 31 March, 2004 on detergents;

•Decision 2006/507/EC of 14 October, 2004 concerning the conclusion, on behalf of the European Community, of the Stockholm Convention on Persistent Organic Pollutants (POPs);

•Directive 2006/7/EC of the European Parliament and of the Council of 15 February, 2006 concerning the management of bathing water quality and repealing Directive 76/160/EEC (with effect from 31 December, 2014);

•Directive 2006/44/EC of 6 September, 2006 on the quality of fresh waters needing protection or improvement in order to support fish life (this Directive will be repealed by the Framework Directive on water as of the end of 2013);

•Directive 2006/11/EC of the European Parliament and of the Council of 15 February, 2006 on pollution caused by certain dangerous substances discharged into the aquatic environment of the Community the European Union (this Directive will be repealed by the Framework Directive on water as of the end of 2013);

•Proposal for a European Parliament and Council Directive of 22 September, 2006 setting out a framework for soil protection and amending Council Directive 2004/35/EC [COM (2006) 231 final – not published in the Official Journal];

•Directive 2006/118/EC of the European Parliament and of the Council of 12 December, 2006 on the protection of groundwater against pollution and deterioration;

2.3.WATER POLICY AND LEGAL FRAMEWORK



•Directive 2007/2/EC of the European Parliament and of the Council of 14 March, 2007 establishing an Infrastructure for Spatial Information in the European Community (INSPIRE);

•Commission Communication of 18 July, 2007: "Addressing the challenge of water scarcity and droughts in the European Union" [COM (2007) 414 final – not published in the Official Journal]; •Directive 2007/60/EC of the European Parliament and of the Council of 23 October, 2007 on the assessment and management of flood risks; aims to manage and reduce the risk of floods, particularly along rivers and in coastal areas;

•Directive 2008/1/EC of the European Parliament and of the Council of 15 January 2008 concerning integrated pollution prevention and control (the IPPC Directive);

•Directive 2009/28/EC on the promotion of the use of energy from renewable sources and amending and subsequently repealing Directives 2001/77/EC and 2003/30/EC.

3. GEOPOLITICAL ASPECTS

Luxembourg is mainly covered by agricultural and forest land. In 2009, 86 per cent of the territory fell in these two categories. Built-up areas occupied about 9 per cent of the country, whereas water and transport infrastructure represented about 5 per cent of the total surface area.

As a member state of the European Union, Luxembourg has to implement this legislation in its own legal framework, defining specific laws for the implementation of the different directives and the common legislative principles. This is a responsibility of the Water Management Administration, which reports to the Minister for the Interior. For example, in 2008, the Water Act was promulgated to consolidate water legislation and transpose the EU Water Framework Directive and Floods Directive into national law.

4. SOURCES

Aquastat – FAO. 1997. Available at: <u>http://www.fao.org/nr/water/aquastat/main/index.stm</u>, [accessed 28 February 2012]

European Environmental Agency (EEA). 2011. The European Environment – State and Outlook 2010. http://www.eea.europa.eu/soer/ [accessed 28 February 2012]